

Boundedness of ϵ -lc Complements on Surfaces

Caucher Birkar^{*†}

February 1, 2008

Contents

1	Boundedness of ϵ-log canonical complements on surfaces	2
1.1	Introduction	2
1.2	Preliminaries	10
1.3	The case of curves	11
1.4	The case of surfaces	12
1.5	Local isomorphic case	13
1.6	Local birational case	20
1.7	Global case	30
1.8	Second proof of the global case	42
1.9	An example	50
1.10	Local cases revisited	51
2	ϵ-log canonical complements in higher dimensions	54
2.1	ϵ -lc complements in dimension 3	56
2.2	ϵ -lc complements in dimension 3: Shokurov's approach	59
3	List of notation and terminology	61
4	References:	62

^{*}Supported by the EPSRC and the Cecil King Prize.

[†]E-mail: caucher.birkar@maths.nott.ac.uk

1 Boundedness of ϵ -log canonical complements on surfaces

1.1 Introduction

The concept of **complement** was introduced and studied by Shokurov [Sh1, Sh2]. He used complements as a tool in the construction of 3-fold log flips [Sh1] and in the classification of singularities and contractions [Sh2]. Roughly speaking a complement is a “good member” of the anti-pluricanonical linear system i.e. a general member of $|-nK_X|$ for some $n > 0$. In order, the existence of such member and the behaviour of the index n are the most important problems in complement theory. Below we give the precise definition of the “good member”.

Throughout this paper we assume that the varieties involved are algebraic varieties over $\mathbb{C}$. In this section the varieties are all surfaces unless otherwise stated. By a log pair $(X/Z, B)$ we mean algebraic varieties X and Z equipped with a projective contraction $X \longrightarrow Z$ and B is an $\mathbb{R}$ -boundary on X . When we write $(X/P \in Z, B)$ we mean a log pair $(X/Z, B)$ with a fixed point $P \in Z$; in such situation we may shrink Z around P without mentioning it. The pair $(X/Z, B)$ is weak log Fano (WLF) if it has log canonical singularities (lc) and $-(K_X + B)$ is nef and big/ Z and X is $\mathbb{Q}$ -factorial.

For the basic definitions of the Log Minimal Model Program (LMMP), the main references are [KMM] and [KM]. And to learn more about the complement theory [Sh2] and [Pr] are the best.

Definition 1.1.1 (Complements) Let $(X/Z, B = \sum_i b_i B_i)$ be a d -dim pair where X is normal and B is an $\mathbb{R}$ -boundary. Then $K_X + B^+$ is called an (ϵ, n) -complement/ $P \in Z$ (resp. in codim 2) for $K_X + B$, where $B^+ = \sum_i b_i^+ B_i$, if the following properties hold:

- ◇ $(X, K_X + B^+)$ is an ϵ -lc pair/ $P \in Z$ (resp. ϵ -lc in codim 2) and $n(K_X + B^+) \sim 0/P \in Z$.
- ◇ $\lfloor (n+1)b_i \rfloor \leq nb_i^+$.

$K_X + B^+$ is called an $(\epsilon, \mathbb{R})$ -complement/ $P \in Z$ (resp. in codim 2) for $K_X + B$ if $(X, K_X + B^+)$ is ϵ -lc/ $P \in Z$ (resp. ϵ -lc in codim 2), $K_X + B^+ \sim_{\mathbb{R}} 0/P \in Z$ and $B^+ \geq B$. An $(\epsilon, \mathbb{Q})$ -complement/ $P \in Z$ can be similarly defined where $\sim_{\mathbb{R}}$ is replaced by $\sim_{\mathbb{Q}}$.

Despite the not quite easy definition above, complements have very good birational and inductive properties which make the theory a powerful tool to apply to the LMMP. In order, complements don't always exist even with strong conditions such as $-(K_X + B)$ being nef [Sh2, 1.1]. But for example they certainly do when $-(K_X + B)$ is nef and big and $K_X + B$ is lc. In this paper, in all the situations that occur the complements usually exist. So we concentrate on the second main problem about complements which is related to several open problems in the LMMP; namely the boundedness. We state conjectures on the boundedness of complements due to Shokurov.

Conjecture 1.1.2 (Weak ϵ -lc Complements) *Let $\Gamma \subseteq [0, 1]$ be a set of real numbers which satisfies D.C.C. Then for any $0 < \delta$ and d there exist a finite set $\mathcal{N}_{\delta,d,\Gamma}$ of positive integers and $0 < \epsilon$ such that any d -dim δ -lc (resp. in codim 2) weak log Fano pair $(X/P \in Z, K_X + B)$, where $B \in \Gamma$, would be (ϵ, n) -complementary/ $P \in Z$ (resp. in codim 2) for some $n \in \mathcal{N}_{\delta,d,\Gamma}$.*

We show the above conjecture as $WC_{\delta,d,\Gamma}$ for short.

Conjecture 1.1.3 (Strong ϵ -lc Complements) *For any $0 < \epsilon$ and d there exists a finite set $\mathcal{N}_{\epsilon,d}$ of positive integers such that any d -dim ϵ -lc (resp. in codim 2) weak log Fano pair $(X/P \in Z, K_X + B)$ has an (ϵ, n) -complement/ $P \in Z$ (resp. in codim 2) for some $n \in \mathcal{N}_{\epsilon,d}$.*

We show the above conjecture as $SC_{\delta,d}$ for short. If we replace $\epsilon > 0$ with $\epsilon = 0$ in the conjecture above (it makes a big difference) then we get the usual conjecture on the boundedness of lc complements which has been studied by Shokurov, Prokhorov and others [Sh2, PSh, PSh1, Pr]. It is proved in dim 2 [Sh2] with some restrictions on the coefficients of B .

The following important conjecture, due to Alexeev and Borisov brothers, is related to the conjectures above [Mc, A1, PSh, MP].

Conjecture 1.1.4 (BAB) *Let $\delta > 0$ be a real number and $\Gamma \subset [0, 1]$. Then those varieties X for which (X, B) is a δ -lc WLF pair of dim d for a boundary $B \in \Gamma$, are elements of an algebraic family.*

We show the above conjecture as $BAB_{\delta,d,\Gamma}$ for short. Alexeev proved $BAB_{\delta,2,\Gamma}$ for any $\delta > 0$ and Γ [A1]. This conjecture was proved by Kawamata for terminal singularities in dim 3 [K1] and $BAB_{1,3,\{0\}}$ was proved by Kollar, Mori, Miyaoka and Takagi [KMMT]. The smooth case was proved by Kollar, Mori and Miyaoka in any dimension. The conjecture is open even in dim 3

when $\delta < 1$. In order, in many interesting applications $\delta < 1$. The following special case of conjecture 1.1.4 was proved by Borisov in dim 3 [B] and by McKernan in any dimension [Mc].

Theorem 1.1.5 (BM) *The set of all klt WLF pairs (X, B) with a fixed given index is bounded i.e. these pairs are elements of an algebraic family.*

The following conjecture is due to Shokurov.

Conjecture 1.1.6 (ACC for mlds) *Suppose $\Gamma \subseteq [0, 1]$ satisfies the descending chain condition (DCC). Then the following set satisfies the ascending chain condition (ACC):*

$$\{\text{mld}(\mu, X, B) | (X, B) \text{ is lc of dim } d, \mu \text{ a point of } X \text{ and } B \in \Gamma\}$$

Note that μ is assumed to be a Grothendieck point of X which is not necessarily closed. We show the above conjecture as $ACC_{d,\Gamma}$ for short. Alexeev proved $ACC_{2,\Gamma}$ for any DCC set $\Gamma \subseteq [0, 1]$ [A2]. This conjecture is open in higher dimensions except in some special cases.

Conjecture 1.1.7 (Log Termination) *Let (X, B) be a Klt pair of dim d . Then any sequence of $K_X + B$ -flips terminates.*

This conjecture is the last step of the LMMP in dim d and we show it as LT_d . Kawamata proved LT_3 [K2] and the four dimensional case with terminal singularities [KMM]. Actually LT_4 is the main missing component of $LMMP_4$ without which we can not apply the powerful LMMP to problems in algebraic geometry. This conjecture did not seem to be that difficult at least because of the short proof of Kawamata to LT_3 where he uses the classification of terminal singularities. The later classification is not known in higher dimensions. Recent attempts by Kawamata and others to solve LT_4 showed that this problem is much deeper than they expected. There is speculation that it may be even more difficult than the flip problem (Shokurov believes this).

We listed several important conjectures with no obvious relation. It is Shokurov's amazing idea to put all these conjectures in a single framework which we call it **Shokurov's Program**:

(1.1.7.1)

ACC_d $\longrightarrow$ LT_d Shokurov proved that the LT_d follows from the above ACC conjecture up to $\dim d$ and the following problem up to $\dim d$ [Sh4]:

Conjecture 1.1.8 (Lower Semi-Continuity) *For any klt pair (X, B) of $\dim d$ and any $c \in \{0, 1, \dots, d-1\}$ the function $\text{mld}_c(\mu, X, B) : \{c - \text{points of } X\} \longrightarrow \mathbb{R}$ is lower semi-continuous.*

A c -point is a c -dim Grothendieck point of X . This conjecture is proved up to $\dim 3$ by Ambro [Am]. This conjecture doesn't seem to be as tough as previous conjectures. Shokurov proved this problem in $\dim 4$ for mlds in $[0, 2]$ [Sh4, lemma 2]. So ACC in $\dim 4$ is enough for the log termination in $\dim 4$ [Sh4, corollary 5]. Actually ACC for mlds in $[0, 1]$ for closed points is enough [Sh4, corollary 5].

BAB_{d-1} $\longrightarrow$ ACC_d Shokurov associates a topological dimension $0 \leq \text{reg}(P \in X, B) \leq d-1$ to any d -dim lc singularity $(P \in X, B)$ [Sh2, 7.9] and proves that the $ACC_{d,\Gamma}$ for pairs with $\text{reg}(P \in X, B) = 0$ is followed from the BAB conjecture in $\dim d-1$ [PSh, 4.4]. In order if $\text{reg}(P \in X, B) = 0$ then the singularity is exceptional (see definition 1.2.8). Also $ACC_{d,\Gamma}$ for pairs with $\text{reg}(P \in X, B) \in \{1, \dots, d-2\}$ can be reduced to lower dimensions. So the only remaining part of $ACC_{d,\Gamma}$ is when $\text{reg}(P \in X, B) = d-1$. This case is expected to be proved using different methods. So *in particular* $ACC_{4,\Gamma}$ follows from the BAB in $\dim 3$ and the $\text{reg}(P \in X, B) = 3$ case. Moreover $ACC_{3,\Gamma}$ is followed from the $\text{reg}(P \in X, B) = 2$ case.

WC_{d-1} $\longrightarrow$ BAB_{d-1} And here it comes probably the most important application of the theory of complements: $WC_{\delta, d-1, \{0\}}$ “implies” $BAB_{\delta, d-1, [0,1]}$. More precisely, these two problems can be solved together at once. In order, in those situations where boundedness of varieties is difficult to prove, boundedness of complements is easier to prove. And that is exactly what we do in this paper for the 2-dim case: we prove $WC_{\delta, 2, \{0\}}$ and $BAB_{\delta, 2, [0,1]}$. Our main objective was to obtain a proof with as little as possible use of surface geometry so that it can be generalised to higher dimensions. In other words, the methods used in the proof of these results are of the most importance to us. After finishing this work, now, we expect to finish the proof of $WC_{\delta, 3, \{0\}}$ “implies” $BAB_{\delta, 3, [0,1]}$ in not a far future! And that was our original goal.

The program in dim 4 Let us mention that by carrying out Shokurov's program in dim 4, in which the main ingredient is $WC_{\delta,3,\{0\}}$ i.e. boundedness of ϵ -lc complements in dim 3, we will prove the following conjectures:

- ACC for mlds in dim 3.
- Boundedness of δ -lc 3-fold log Fanos i.e. BAB in dim 3.
- ACC for mlds in dim 4.
- Lower semi-continuity for mlds in dim 4.
- Log termination in dim 4 and then LMMP in dim 4.

About this paper:

1. Section 1 is devoted to the study of complements on log surfaces.
2. In 1.2 we recall some definitions and lemmas.
3. In 1.3 we prove $WC_{\delta,1,[0,1]}$ i.e. the boundedness of ϵ -lc complements in dim 1 (**theorem 1.3.1**).
4. In 1.5 we prove $WC_{\delta,2,\{0\}}$ for the case $X = Z$ i.e. the boundedness of ϵ -lc complements in dim 2, locally, for points on surfaces with $B = 0$ (**theorem 1.5.1**).
5. In 1.6 we prove $WC_{\delta,2,\{0\}}$ when X/Z is a birational equivalence i.e. the boundedness of ϵ -lc complements in dim 2, locally, for birational contractions of surfaces with $B = 0$ (**theorem 1.6.1**). This proof is a surface proof i.e. we heavily use surface geometry and it is not expected to be generalized to higher dimensions. A second proof of the birational case is given in 1.10 (**theorem 1.10.1**).
6. In 1.7 we prove $WC_{\delta,2,\{0\}}$ when $Z = pt$ i.e. the boundedness of ϵ -lc complements on surfaces, globally, with $B = 0$ (**theorem 1.7.1**). The proof is based on the LMMP and we expect to be generalised to higher dimensions. As a corollary we give a totally new proof to the boundedness of ϵ -lc log Dell Pezzo surfaces (=BAB indim 2) (**corollary 1.7.9**). Another application of our theorem is that the boundedness of lc ($\epsilon = 0$) complements can be proved only using complement theory (**theorem 1.1.15**). The later boundedness was proved by Shokurov [Sh2].

7. In 1.8 we give a second proof of $WC_{\delta,2,\{0\}}$ in the global case i.e. when $Z = pt$ (**theorem 1.8.1**). This proof also uses surface geometry and is not expected to be generalized to higher dimensions.
8. In 1.9 we discuss an example which shows that the transformed boundary doesn't have a better singularity than the original boundary.
9. In 1.10 we give a proof to all local cases for $B \in \Phi_{sm}$, in particular, the case where X/Z is a fibration over a curve (**theorem 1.10.1**). This proof is also based on the LMMP.
10. Section 2 is about higher dimensional ϵ -lc complements. We discuss our joint work with Shokurov.
11. In 2.1 We give a **Plan** about how to attack the boundedness of ϵ -lc complements in dimension 3. This is proposed by the author.
12. In 2.2 We give Shokurov's **Plan** about how to attack the boundedness of ϵ -lc complements in dimension 3.

Let us summarise the main results of section 1:

Theorem 1.1.9 *Conjecture 1.1.2 holds in dim 1 for $\Gamma = [0, 1]$.*

See 1.3.1 for the proof.

Theorem 1.1.10 *Conjecture 1.1.2 holds in dim 2 in the global case (i.e. $\dim Z = 0$) for $\Gamma = \{0\}$.*

See 1.7.1 and 1.8.1 for proofs.

Theorem 1.1.11 *Conjecture 1.1.2 holds in dim 2 in the local cases (i.e. $\dim Z > 0$) for $\Gamma = \Phi_{sm}$.*

See 1.10.1 , 1.5.1 and 1.6.1 for proofs.

Corollary 1.1.12 *Conjecture 1.1.4 holds in dim 2.*

See 1.7.9 for proof. Conjecture 1.1.4 in dim 2 was first proved by Alexeev using different methods [A1].

Corollary 1.1.13 *Theorem 1.1.15 can be proved using only the complement theory.*

See the discussion following theorem 1.1.15.

Remark 1.1.14 (ϵ -lc Complements Method) Though formally speaking the list above are the main results in section 1, but we believe that the method used to prove 1.7.1 and 1.10.1 is the most important result of this section.

Here we mention some developments in the theory of complements. The following theorem was proved by Shokurov [Sh2] for surfaces.

Theorem 1.1.15 *There exists a finite set $\mathcal{N}_2$ of positive integers such that any 2-dim lc weak log Fano pair $(X/P \in Z, B)$ has a $(0, n)$ -complement/ $P \in Z$ for some $n \in \mathcal{N}_2$ if B is semi-standard i.e. for each coefficient b of B , $b \geq \frac{6}{7}$ or $b = \frac{m-1}{m}$ for some natural number m . In order if $\dim Z > 0$ then the theorem holds for a general boundary.*

In order Shokurov uses the BAB in dim 2 in the proof of the above theorem. As mentioned before, the results of this paper imply the BAB in dim 2. So the above theorem can be proved only based on the theory of complements. A similar theorem is proved by Prokhorov and Shokurov in dim 3 modulo BAB in dim 3 and the effective adjunction in dim 3 [PSh1]. However the local case doesn't need the later assumptions as the following theorem shows.

Theorem 1.1.16 *Let $(X/P \in Z, B)$ be a Klt WLF 3-fold pair where $\dim Z \geq 1$ and $B \in \Phi_{sm}$. Then $K_X + B$ is $(0, n)$ -complementary/ $P \in Z$ for some $n \in \mathcal{N}_2$.*

Complements have good inductive properties as the theorem above shows which was proved by Prokhorov and Shokurov [PSh]. This theorem is stated and proved in higher dimensions in more general settings (see [PSh]). To avoid some exotic definitions, we stated only the 3-fold version.

Finally we give some easy examples of complements. More interesting examples can be found in [Sh1, Sh2, Pr, PSh, PSh1].

Example 1.1.17 Let $(X/Z, B) = (\mathbb{P}^1/pt., 0)$ and P_1, P_2, P_3 distinct points on $\mathbb{P}^1$. Then $K_X + P_1 + P_2$ is a $(0, 1)$ -complement for K_X but it is not an (ϵ, n) -complement for any $\epsilon > 0$ since $K_X + P_1 + P_2$ is not Klt. On the other hand $K_X + \frac{2}{3}P_1 + \frac{2}{3}P_2 + \frac{2}{3}P_3$ is a $(\frac{1}{3}, 3)$ -complement for K_X .

Example 1.1.18 Let $(X_1/Z_1, B_1) = (\mathbb{P}^2/pt., 0)$ and $(X_2/Z_2, B_2) = (\mathbb{P}^2/\mathbb{P}^2, 0)$. Then K_{X_2} is a $(2, 1)$ -complement/ Z_2 at any point $P \in Z_2$ but obviously K_{X_1} is not even numerically zero/ Z_1 though $K_{X_1} = K_{X_2}$.

Example 1.1.19 Let $(X/Z, B) = (X/X, 0)$ where X is a surface with canonical singularities. Then the index of K_X is 1 at any point $P \in X$. So we can take $B^+ = 0$ and K_X a $(1, 1)$ -complement/ X for K_X at any $P \in X$.

acknowledgement

I am deeply grateful to my adviser Professor V.V. Shokurov for introducing me to these problems and helping me through very patiently. I also thanks the EPSRC and the Cecil King Prize for their financial support.

1.2 Preliminaries

In this subsection we bring some basic definitions and constructions.

Definition 1.2.1 A set $\mathcal{X}$ of varieties of the same dimension is called bounded if there are schemes $\mathbb{X}$ and S of finite type and a morphism $\phi : \mathbb{X} \rightarrow S$ such that each element of $\mathcal{X}$ is isomorphic to a geometric fibre of ϕ . Moreover each fibre should give an element of $\mathcal{X}$.

Definition 1.2.2 Let $\mathcal{X}$ be a set of pairs (X, B_X) of the same dimension. Then the set consisting of all $(X, \text{Supp } B_X)$ where $X \in \mathcal{X}$ is called bounded if there are schemes $\mathbb{X}$ and S of finite type, a divisor $\mathbb{B}$ on $\mathbb{X}$ and a morphism $\phi : \mathbb{X} \rightarrow S$ such that each $X \in \mathcal{X}$ is isomorphic to a geometric fibre $\mathbb{X}_s$ and $\text{Supp } B_X = \text{Supp } \mathbb{B}|_{\mathbb{X}_s}$. And (X, B) bounded means that $(X, \text{Supp } B_X)$ is bounded and there are only finitely many possibilities for the coefficients of B .

Definition 1.2.3 Let (X, B) be a Klt pair of dim d . Let $\phi : Y \rightarrow X$ be a morphism such that $B_Y \geq 0$ where $K_Y + B_Y = \phi^*(K_X + B)$. Then Y is called a partial resolution of (X, B) .

Lemma 1.2.4 Let $\mathcal{X} = \{X\}$ be a bounded set of Klt varieties of dim d such that $-K_X$ is nef and big. Then $\mathcal{Y}$ the set of partial resolutions for all $X \in \mathcal{X}$ is bounded.

Proof By assumptions for each $Y \in \mathcal{Y}$ there are $X \in \mathcal{X}$ and a boundary B_Y such that $K_Y + B_Y = \phi^*K_X$. Since $\mathcal{X}$ is bounded hence there are only a finite number of possibilities for the coefficients of B_Y independent of Y thus the index of $K_Y + B_Y$ is bounded. And since $-K_X$ is nef and big so $-(K_Y + B_Y)$ is also nef and big. Now by Borisov-Mckernan [Mc] the set $\mathcal{Y}$ is bounded. $\square$

Definition 1.2.5 A variety X/Z of dim d , is called Pseudo-WLF/ Z if there exists a boundary B such that $(X/Z, B)$ is WLF. Moreover X is called Klt Pseudo-WLF/ Z if there is a Klt WLF $(X/Z, B)$. If $\dim Z = 0$ then we usually drop Z .

Remark 1.2.6 Pseudo-WLF varieties have good properties. For example $\overline{NE}(X/Z)$ is a finite rational polyhedral cone. Moreover each extremal face

of the cone is contractible [Sh5, Sh3]. In section 2 we prove that the Klt Pseudo-WLF property is preserved under flip and divisorial contractions.

Definition 1.2.7 The set $\Phi_{sm} = \{\frac{k-1}{k} | k \in \mathbb{N}\} \cup \{1\}$ is called the set of standard boundary multiplicities. For a boundary B by $B \in \Phi_{sm}$ we mean that the coefficients of B are in Φ_{sm} .

Definition 1.2.8 (Exceptional pairs) Let $(X/Z, B)$ be a pair of dim d . If $\dim Z = 0$ then $(X/Z, B)$ is called exceptional if there is at least a $(0, \mathbb{Q})$ -complement $K_X + B^+$ and any $(0, \mathbb{Q})$ -complement $K_X + B^+$ is Klt. If $\dim Z > 0$ then $(X/Z, B)$ is called exceptional if there is at least a $(0, \mathbb{Q})$ -complement $K_X + B^+$ and any $(0, \mathbb{Q})$ -complement $K_X + B^+$ is plt on a log terminal resolution. Otherwise $(X/Z, B)$ is called non-exceptional.

Remark 1.2.9 Boundedness of analytic (ϵ, n) -complements implies the boundedness of algebraic (ϵ, n) -complement. That is because of the general GAGA pricipile [Sh1].

Lemma 1.2.10 *Let $Y/X/Z$ and $K_Y + B_Y$ be nef/ X and $K_X + B = *(K_Y + B_Y)$ be (ϵ, n) -complementary/ Z . Moreover assume that each non-exceptional/ X component of B_Y that intersects an exceptional divisor/ X has a standard coefficient then (Y, B_Y) will also be (ϵ, n) -complementary/ Z .*

Proof See [PSh, 6.1].

1.3 The case of curves

In this subsection we prove 1.1.2 for the case of curves. Note that 1-dim global log Fano pairs are just $(\mathbb{P}^1, B)$ for a boundary $B = \sum_i b_i B_i$ where $\sum_i b_i - 2 < 0$. The local case for curves is trivial.

Theorem 1.3.1 *$WC_{\delta, 1, [0, 1]}$ holds; more precisely, suppose $\frac{m-1}{m} \leq 1 - \delta < \frac{m}{m+1}$ for m a natural number then we have:*

- ◇ $N_{\delta, 1, [0, 1]} \subseteq \cup_{0 < k \leq m} \{k, k+1\}$.
- ◇ $(\mathbb{P}^1, B^+)$ can be taken $\frac{1}{m+1}$ -lc.

Proof Let $B = \sum_i b_i B_i$ and put $b = b_h = \max\{b_i\}$ and suppose $\frac{k-1}{k} \leq b < \frac{k}{k+1}$ for a natural number k . If $k = 1$ then $0 \leq b < \frac{1}{2}$ and so we have a 1-complement $B^+ = 0$.

Now assume that $K > 1$ and define $a_{i,t} = \lfloor (t+1)b_i \rfloor$ and note that by our assumptions $\sum_i a_{i,k} \leq 2k+1$ since $\sum_i b_i < 2$. If $K+B$ doesn't have a k -complement then $\sum_i a_{i,k} = 2k+1$. Since $\frac{k-1}{k} \leq b < \frac{k}{k+1}$ we have $\frac{(k+1)(k-1)}{k} = k+1 - \frac{k+1}{k} = k - \frac{1}{k} \leq (k+1)b < \frac{(k+1)k}{k+1} = k$. Thus $a_{h,k} = k-1$ and $1 - \frac{1}{k} \leq \langle (k+1)b \rangle < 1$ where $\langle \cdot \rangle$ stands for the fractional part.

Now $a_{i,k+1} = \lfloor (k+2)b_i \rfloor = \lfloor (k+1)b_i + b_i \rfloor$. So $a_{i,k+1}$ is $a_{i,k}$ or $a_{i,k} + 1$. The later happens iff $1 \leq b_i + \langle (k+1)b_i \rangle$. By the above $b_h + \langle (k+1)b_h \rangle \geq \frac{k-1}{k} + 1 - \frac{1}{k} \geq 1$ so $a_{h,k+1} = a_{h,k} + 1$. On the other hand since $\sum_i b_i < 2$ and $\sum_i a_{i,k} = 2k+1$ then $\sum_i \langle (k+1)b_i \rangle < 1$. And since $1 - \frac{1}{k} \leq \langle (k+1)b_h \rangle$ then $\langle (k+1)b_i \rangle < \frac{1}{k}$ if $i \neq h$. So under assumption $i \neq h$ if $1 \leq \langle (k+1)b_i \rangle + b_i$ then $1 - \frac{1}{k} < b_i$.

Thus if $K+B$ has no $k+1$ -complement then $1 \leq \langle (k+1)b_j \rangle + b_j$ should hold at least for some $j \neq h$. So again we have $1 - \frac{1}{k} \leq \langle (k+1)b_j \rangle$ and so $\langle (k+1)b_j \rangle + \langle (k+1)b_h \rangle \geq 2(1 - \frac{1}{k}) \geq 1$ and this is a contradiction. Hence $K+B$ should have a k or $k+1$ -complement. If $K+B$ has a k complement then we can have a maximum $\max\{b_i^+\} = b^+ = 1 - \frac{1}{k} \leq 1 - \delta$. If it has a $k+1$ -complement then again we can have a maximum $b^+ \leq \frac{k}{k+1}$. Since $0 < k \leq m$ then $N_{\delta,1} \subseteq \cup_{0 < k \leq m} \{k, k+1\}$ and $K+B^+$ can be chosen as $\frac{1}{m+1}$ -lc. These prove the theorem. $\square$

Remark 1.3.2 Above we just proved that $\sum_i \frac{\lfloor (n+1)b_i \rfloor}{n} \leq 2$ for a bounded n . If the equality doesn't hold then we may add some positive coefficients to get the equality and construct the complement.

1.4 The case of surfaces

We divide the surface case of conjecture 1.1.2 into the following cases:

local isomorphic X/Z is an isomorphism.

local birational X/Z is birational but may not be an isomorphism.

local over curve Z is a curve.

global Z is a point.

1.5 Local isomorphic case

The main theorem in this subsection is theorem 1.5.1. We use classification of surface singularities.

Theorem 1.5.1 *Conjecture $WC_{\delta,2,\{0\}}$ holds in the local isomorphic case.*

Proof If $\delta \geq 1$ then P is smooth and so we are already done. So assume that $\delta < 1$. If the singularity at P is of type E_6 , E_7 or E_8 then there are only a finite number of possibilities up to analytic isomorphism because of the δ -lc assumption [see Pr 6.1.2]. Otherwise the graph of the resolution will be either of type A_r :

$$O^{-\alpha_r} \text{ --- } \dots \text{ --- } O^{-\alpha_2} \text{ --- } O^{-\alpha_1}$$

where $\alpha_i \geq 2$. Or of type D_r :

$$\begin{array}{c} O^{-2} \\ | \\ O^{-\alpha_r} \text{ --- } \dots \text{ --- } O^{-\alpha_2} \text{ --- } O^{-\alpha_1} \\ | \\ O^{-2} \end{array}$$

where $\alpha_i \geq 2$.

First we work out the case A_n . Let $K_W - \sum_i e_i E_i = {}^*K_Z$ where e_i are the discrepancies for a log resolution $W \rightarrow Z/P$. The following lemma is well known and a proof can be found in [AM, 1.2].

Lemma 1.5.2 *The numbers $(-E_i^2)$ are bounded from above in terms of δ .*

Intersecting $K_W - \sum_i e_i E_i$ with all the exceptional divisors we get a system like the following:

$$\begin{aligned} a_1(-E_1^2) - a_2 - 1 &= 0 \\ a_2(-E_2^2) - a_1 - a_3 &= 0 \\ a_3(-E_3^2) - a_2 - a_4 &= 0 \end{aligned}$$

$$\begin{aligned}
& \vdots \\
& a_{r-1}(-E_{r-1}^2) - a_{r-2} - a_r = 0 \\
& a_r(-E_r^2) - a_{r-1} - 1 = 0
\end{aligned}$$

where a_i is the log discrepancy of E_i with respect to K_Z .

From the equation $a_i(-E_i^2) - a_{i-1} - a_{i+1} \leq 0$ we get $a_i(-E_i^2 - 2) + a_i - a_{i-1} \leq a_{i+1} - a_i$ which shows that if $a_{i-1} \leq a_i$ then $a_i \leq a_{i+1}$ and moreover if $a_{i-1} < a_i$ then $a_i < a_{i+1}$. So the solution for the system above should satisfy the following:

$$(1.5.2.1)$$

$$a_1 \geq \cdots \geq a_i \leq \cdots \leq a_r$$

for some $i \geq 1$. If $r \leq 2$ (or any fixed number) then the theorem is trivial. So we may assume that $r > 3$ and also can assume $i \neq r$ unless $a_1 = a_2 = \cdots = a_r$. Now for any $i \leq j < r$ if $-E_j^2 > 2$ then $a_{j+1} - a_j \geq a_j(-E_j^2 - 2) \geq \delta$. So if we have l members in $\{j : -E_j^2 > 2 \text{ and } i \leq j < r\}$ then $a_r \geq l\delta$. Hence $a_r(-E_r^2 - 1) + a_r - a_{r-1} \geq l\delta$ which contradicts the last equation in the system if l gets big arbitrarily. In order $l\delta \leq 1$ and so $l \leq \frac{1}{\delta}$. Similar observation shows that the number l' of $1 \leq j \leq i$ where $-E_j^2 > 2$ should be bounded. Then $l + l' \leq \frac{2}{\delta}$.

Suppose $a_{i_2} = \cdots = a_i = \cdots = a_{i_1}$, $a_{i_1-1} \neq a_{i_1}$ (or $i_1 = 1$) and $a_{i_2} \neq a_{i_2+1}$ (or $i_2 = r$) where $i_2 \leq i \leq i_1$. Assume that $i_1 \neq i$ or $i_2 \neq i$. Let for example $i_1 \neq i$. Then if all a_j are not equal ($= 1$) then we have

$$\begin{aligned}
1 &= (-E_r^2 - 1)a_r + a_r - a_{r-1} \geq (r - i_1)(a_{i_1+1} - a_{i_1}) \\
&= (r - i_1)[(-E_{i_1}^2 - 2)a_{i_1} + a_{i_1} - a_{i_1-1}] \\
&= (r - i_1)(-E_{i_1}^2 - 2)a_{i_1} \geq (r - i_1)\delta
\end{aligned}$$

because $-E_{i_1}^2$ can not be equal to 2.

So $(r - i_1)\delta \leq 1$ then $r - i_1 \leq \frac{1}{\delta}$ is bounded. Similarly i_2 should be bounded. These observation show that, mentioning that $-E_k^2$ are bounded, the denominators of a_k are bounded. And so the index of K_Z at P is bounded and so we are done in this case. But if $i_1 = i = i_2$ then the story is different. In this case note that $\delta \leq (-E_i^2 - 2)a_i = a_{i-1} - a_i + a_{i+1} - a_i$. So $\frac{\delta}{2} \leq a_{i-1} - a_i$ or $\frac{\delta}{2} \leq a_{i+1} - a_i$. For example assume that the later holds then similar to

the calculations we just carried out above, $r - i$ will be bounded. But it can happen that $a_{i-1} - a_i$ is very small so we won't be able to bound i .

In order, we try to find a solution for the following system with bounded denominators:

$$\begin{aligned}
u_1(-E_1^2) - u_2 - 1 &\leq 0 \\
u_2(-E_2^2) - u_1 - u_3 &\leq 0 \\
u_3(-E_3^2) - u_2 - u_4 &\leq 0 \\
&\vdots \\
u_{r-1}(-E_{r-1}^2) - u_{r-2} - u_r &\leq 0 \\
u_r(-E_r^2) - u_{r-1} - 1 &\leq 0
\end{aligned}$$

To do this, note that if $-E_{i-1}^2 > 2$ then $\delta \leq (-E_{i-1}^2 - 2)a_{i-1} = a_{i-2} - a_{i-1} + a_i - a_{i-1} \leq a_{i-2} - a_{i-1}$ then again similar computations to the above shows that i is bounded. Now let j be the smallest number such that $-E_j^2 = \dots = -E_{i-1}^2 = 2$ (remember that we have assumed $\frac{\delta}{2} \leq a_{i+1} - a_i$). Hence j is bounded. Now take $u_j = \dots = u_i = \frac{1}{2}$ then the following equations are satisfied if $i - j > 2$:

$$\begin{aligned}
u_{j+1}(-E_{j+1}^2) - u_j - u_{j+2} &= 2u_j - u_j - u_j = 0 \\
&\vdots \\
u_{i-1}(-E_{i-1}^2) - u_{i-2} - u_i &= 2u_{i-1} - u_{i-2} - u_i = 0
\end{aligned}$$

Since $r - i$ and j are bounded then the number of remaining equations is bounded and so to satisfy them we just have to divide $u_i = \frac{1}{2}$ by a bounded natural number. This finishes the A_r type. The D_r type will follow.

Remark 1.5.3 In order we have constructed a Klt log divisor $K_W + D$ with bounded index such that $-(K_W + D)$ is nef and $\text{big}/P \in Z$. Now we may use remark 1.2.9.

Remark 1.5.4 All the bounds occurring in the proof are effective and can be calculated in terms of δ .

Remark 1.5.5 In Shokurov's case where $\delta = \epsilon = 0$ we just take $u_1 = \dots = u_r = 0$.

The case of D_r : We have a chain $E_1, \dots, E_r$ of exceptional divisors plus E and E' where E intersects only E_1 and the same holds for E' . In this case we have the following system:

$$\begin{aligned}
a(-E^2) - a_1 - 1 &= 0 \\
a'(-E'^2) - a_1 - 1 &= 0 \\
a_1(-E_1^2) - a - a' - a_2 + 1 &= 0 \\
a_2(-E_2^2) - a_1 - a_3 &= 0 \\
a_3(-E_3^2) - a_2 - a_4 &= 0 \\
&\vdots \\
a_{r-1}(-E_{r-1}^2) - a_{r-2} - a_r &= 0 \\
a_r(-E_r^2) - a_{r-1} - 1 &= 0
\end{aligned}$$

Note that $-E^2 = -E'^2 = 2$ so $2a - a_1 - 1 = 0$ and $2a' - a_1 - 1 = 0$ hence $a + a' = a_1 + 1$. Replacing this in the third equation and ignoring the two first equations we get the following system:

$$\begin{aligned}
a_1(-E_1^2 - 1) - a_2 &= 0 \\
a_2(-E_2^2) - a_1 - a_3 &= 0 \\
a_3(-E_3^2) - a_2 - a_4 &= 0 \\
&\vdots \\
a_{r-1}(-E_{r-1}^2) - a_{r-2} - a_r &= 0 \\
a_r(-E_r^2) - a_{r-1} - 1 &= 0
\end{aligned}$$

From this system we get a solution as following:

$$a_1 = \dots = a_i < a_{i+1} < \dots < a_r$$

($i=r$ also may happen. In this case $a = a' = a_1 = \dots = a_r = 1$). Now $r - i$ should be bounded. In order if $i > 1$ then $-E_1^2 = \dots = -E_{i-1}^2 = 2$ but $-E_i^2 > 2$ (we have assumed $r > i$). Now $\delta(-E_i^2 - 2) \leq a_i(-E_i^2 - 2) + a_i - a_{i-1} = a_{i+1} - a_i$ (if $i = 1$ then $\delta(-E_1^2 - 2) \leq a_1(-E_1^2 - 2) = a_2 - a_1$). We also have the fact that $a_{k+1} - a_k \leq a_{k+2} - a_{k+1}$ for $i \leq k < r - 1$. And on the other hand $\sum_{i \leq k < r} a_{k+1} - a_k \leq a_r < a_r + a_r - a_{r-1} < 1$. So we conclude

that $r - i$ should be bounded. Moreover since $-E_k^2$ is bounded, this proves that the denominators of all a_k in the D_r case are bounded and so the index of K_Z at P . In this case $B^+ = 0$ and this finishes the proof of theorem 1.5.1. $\square$

Remark 1.5.6 Essentially the boundedness properties that we proved and used in the proof of theorem 1.5.1 had been more or less discovered by some other people independently. Shokurov had used these ideas in an unpublished preprint on mlds.

Recall 1.5.7 Here we recall the diagrams for the E_6 , E_7 and E_8 types of singularities. The following is a general case of such singularities:

$$\begin{array}{ccccc} \mathbb{C}^2/\mathbb{Z}_{m_1} & \text{---} & O^{-p} & \text{---} & \mathbb{C}^2/\mathbb{Z}_{m_2} \\ & & \downarrow & & \\ & & O^{-2} & & \end{array}$$

where the only possibilities for (m_1, m_2) are $(3, 3)$, $(3, 4)$ and $(3, 5)$. So the possible diagrams are as follows: For $(m_1, m_2) = (3, 3)$ we have

1

$$\begin{array}{ccccc} O^{-3} & \text{---} & O^{-p} & \text{---} & O^{-3} \\ & & \downarrow & & \\ & & O^{-2} & & \end{array}$$

2

$$\begin{array}{ccccccc} O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-p} & \text{---} & O^{-3} \\ & & & & \downarrow & & \\ & & & & O^{-2} & & \end{array}$$

3

$$\begin{array}{ccccccccc} O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-p} & \text{---} & O^{-2} & \text{---} & O^{-2} \\ & & & & \downarrow & & & & \\ & & & & O^{-2} & & & & \end{array}$$

For $(m_1, m_2) = (3, 4)$ we have

4

$$\begin{array}{ccccc} O^{-3} & \text{---} & O^{-p} & \text{---} & O^{-4} \\ & & | & & \\ & & O^{-2} & & \end{array}$$

5

$$\begin{array}{ccccc} O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-p} & \text{---} & O^{-4} \\ & & & & | & & \\ & & & & O^{-2} & & \end{array}$$

6

$$\begin{array}{ccccccc} O^{-3} & \text{---} & O^{-p} & \text{---} & O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-2} \\ & & | & & & & & & \\ & & O^{-2} & & & & & & \end{array}$$

7

$$\begin{array}{ccccccc} O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-p} & \text{---} & O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-2} \\ & & & & | & & & & & & \\ & & & & O^{-2} & & & & & & \end{array}$$

And for $(m_1, m_2) = (3, 5)$ we have

8

$$\begin{array}{ccccc} O^{-3} & \text{---} & O^{-p} & \text{---} & O^{-5} \\ & & | & & \\ & & O^{-2} & & \end{array}$$

9

$$\begin{array}{ccccc} O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-p} & \text{---} & O^{-5} \\ & & & & | & & \\ & & & & O^{-2} & & \end{array}$$

10

$$\begin{array}{ccccc} O^{-3} & \text{---} & O^{-p} & \text{---} & O^{-2} & \text{---} & O^{-3} \\ & & | & & & & \\ & & O^{-2} & & & & \end{array}$$

11

$$\begin{array}{ccccccc} O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-p} & \text{---} & O^{-2} & \text{---} & O^{-3} \\ & & & & | & & & & \\ & & & & O^{-2} & & & & \end{array}$$

12

$$\begin{array}{ccccccc} O^{-3} & \text{---} & O^{-p} & \text{---} & O^{-3} & \text{---} & O^{-2} \\ & & | & & & & \\ & & O^{-2} & & & & \end{array}$$

13

$$\begin{array}{ccccccc} O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-p} & \text{---} & O^{-3} & \text{---} & O^{-2} \\ & & & & | & & & & \\ & & & & O^{-2} & & & & \end{array}$$

14

$$\begin{array}{ccccccc} O^{-3} & \text{---} & O^{-p} & \text{---} & O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-2} \\ & & | & & & & & & \\ & & O^{-2} & & & & & & \end{array}$$

15

$$\begin{array}{ccccccc} O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-p} & \text{---} & O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-2} & \text{---} & O^{-2} \\ & & & & | & & & & \\ & & & & O^{-2} & & & & \end{array}$$

1.6 Local birational case

In this subsection wherever we write $/Z$ we mean $/P \in Z$ for a fixed point P on Z .

Theorem 1.6.1 $WC_{\delta,2,\{0\}}$ holds in the birational case.

Strategy of the proof: Let W be a minimal resolution of X and $\{E_i\}$, $\{F_j\}$ be the exceptional divisors $/Z$ on W where the E_i are exceptional $/X$ but F_j are not (E will be used for a typical E_i and similarly F for F_j or its birational transform). We will construct an antinef $/Z$ and Klt log divisor $K_W + \Omega = K_W + \sum_i u_i E_i + \sum_j u_j F_j$ where $u_i, u_j < 1$ are rational numbers with bounded denominators. Then we use remark 1.2.9.

Proof By contracting those curves where $-K_X$ is numerically zero, we can assume that $-K_X$ is ample $/Z$ (we can pull back the complement). Let W be the minimal resolution of X . Then since K_W is nef $/X$ by the negativity lemma we have $K_W - \sum_i e_i E_i = K_W + \sum_i (1 - e_i) E_i \equiv {}^*K_X$ where $e_i \leq 0$.

Definition 1.6.2 For any smooth model Y where $W/Y/Z$ we define $\overline{exc}(Y/Z)$ to be the graph of the exceptional curves ignoring the birational transform of exceptional divisors of type F . For an exceptional $/Z$ divisor G on Y not of type F , $\overline{exc}(Y/Z)_G$ means the connected component of $\overline{exc}(Y/Z)$ where G belongs to.

Lemma 1.6.3 We have the followings on W :

- ◇ The exceptional divisors $/Z$ on W are with simple normal crossings.
- ◇ Each F (i.e. each exceptional divisor of type F) is a -1 -curve.
- ◇ The model $\overline{W}$ obtained by blowing down -1 -curves $/Z$ is the minimal resolution of Z .
- ◇ Each F cuts at most two exceptional divisors of type E .

Proof Let F be an exceptional divisor $/Z$ on W which is not exceptional $/X$. Then $(K_W - \sum_i e_i E_i) \cdot F = K_W \cdot F + \sum_i (-e_i) E_i \cdot F = 2p_a(F) - 2 - F^2 + \sum_i (-e_i) E_i \cdot F < 0$ where $p_a(F)$ stands for the arithmetic genus of the curve F . Then $2p_a(F) - 2 - F^2 < 0$ and so $p_a(F) = 0$ and $-F^2 = 1$. In other

words F is a -1 -curve.

On the other hand by contracting -1 -curves/ Z (i.e. running the classical minimal model theory for smooth surfaces on W/Z) we get a model $\overline{W}/Z$ where $K_{\overline{W}}$ is nef/ Z . Actually $\overline{W}$ is the minimal resolution of $P \in Z$.

The exceptional divisors/ Z on $\overline{W}$ are with simple normal crossings and since W is obtained from $\overline{W}$ by a sequence of blow ups then the exceptional divisors/ Z on W also would be with simple normal crossings. This observation gives more information. Since all the F , exceptional/ Z but not/ X , are contracted/ $\overline{W}$ then they should intersect at most two of E_i because $exc(\overline{W}/Z)$ is with simple normal crossings and F is exceptional/ $\overline{W}$. Moreover no two exceptional divisors of type F should intersect on W because they are both -1 -curves. This means that the intersection points of any two exceptional divisor/ Z on X should be a singular point of X . Also any exceptional divisor/ Z on X contains at most two singular points of X . $\square$

Now Suppose $\{Q_k\}_k$ to be the singular points of X . If no one of the points $\{Q_k\}$ is of type A_r then the proof of theorem 1.5.1 shows that the discrepancies e_i are with bounded denominators so we are already done. But if there is one point of type A_r then the proof is more complicated (surprisingly the A_r type is the most simple case in the sense of Shokurov i.e. when $\delta = 0$). Similar to the proof of theorem 1.5.1 we try to understand the structure of $exc(W/Z)$ and the blow ups $W \rightarrow \overline{W}$.

Definition 1.6.4 A smooth model $\check{W}$ where $W/\check{W}$ and $\check{W}/\overline{W}$ are series of smooth blow ups, is called a *blow up model* of $\overline{W}$. Such a model is called *minimal* if there is X' such that $K_{\check{W}}$ is nef/ X' and $X/X'/Z$. In other words it is the minimal resolution of X' . The connected components of $\overline{exc}(\check{W}/Z)$ are either of type A_r , D_r , E_6 , E_7 or E_8 for a minimal blow up model.

Definition 1.6.5 We call the divisor $K_W + \omega = K_W + \sum_i (1 - a_i) E_i = {}^*K_X$ the *primary log divisor*. The pair (X, B) has a log canonical n -complement $K_X + B^+$ over Z in the sense of Shokurov [Sh2] ($n \leq 6$). From now on we call it a *Shokurov complement*. So $K_W + \omega_{Sh} + C = K_W + \sum_i (1 - a_i^{Sh}) E_i + \sum_j (1 - a_j^{Sh}) F_j + C = {}^*(K_X + B^+)$ where C is the birational transform of the non-exceptional part of B^+ . We call $K_W + \omega_{Sh}$ a *Shokurov log divisor* and the numbers a_i^{Sh} and a_j^{Sh} Shokurov log discrepancies.

Definition 1.6.6 In the graph $exc(W/Z)$ if we ignore those F which appear with zero coefficient in ω_{Sh} (i.e. $a^{Sh} = 1$) then we get a graph

$exc(W/Z)_{>0}$ with some connected components. The connected graph $\mathcal{C}$ consisting of exceptional/ Z curves with $a^{Sh} = 0$, is in one of the components of the graph $exc(W/Z)_{>0}$ which we show by $\mathcal{G}$ ($\mathcal{C}$ is connected because of the connectedness of the locus of log canonical centres/ $P \in Z$). Now contracting all -1 -curves/ Z in $\mathcal{G}$ and continuing the contractions of subsequent -1 -curves/ Z which appear in $\mathcal{G}$, finally we get a model which we show as $W_{\mathcal{G}}$. The transform of $\mathcal{G}$ on $W_{\mathcal{G}}$ is shown by $\mathcal{G}_1$ and similarly the transform of $\mathcal{C}$ is $\mathcal{C}_1$.

Definition 1.6.7 A chain of exceptional curves consisting of $G_{\beta_1}, \dots, G_{\beta_r}$ is called *strictly monotonic* if $r = 1$ or if $a_{\beta_1} < a_{\beta_2} < \dots < a_{\beta_r}$ (these are log discrepancies with respect to K_X). G_{β_1} is called the *base curve*.

Definition 1.6.8 Let $G \in exc(\ddot{W}/Z)$ for a smooth blow up model $\ddot{W}$. Then define the *negativity* of G on this model as $N_{\ddot{W}}(G) = (K_{\ddot{W}} + *_\omega).G \leq 0$ ($*_\omega$ is pushdown of ω). We also define the *total negativity* by $N_{\ddot{W}} = \sum_{\alpha} N_{\ddot{W}}(G_{\alpha})$ where G_{α} runs on all exceptional divisors/ Z on $\ddot{W}$ and for $G \in \overline{exc}(\ddot{W}/Z)$ define $N_{\ddot{W},G} = \sum_{\alpha} N_{\ddot{W}}(G_{\alpha})$ when the sum runs on all members of $\overline{exc}(\ddot{W}/Z)_G$. Similarly define the negativity functions N^{Sh} and N^+ replacing ω with ω_{Sh} and $\omega_{Sh} + C$ respectively. Note that the later is always zero, because $K_W + \omega_{Sh} + C \equiv 0/Z$.

Definition 1.6.9 The (smooth) blow up of a point which belongs to two exceptional divisors/ Z on a model is called a *double blow up*. If this point belongs to just one exceptional divisor/ Z then we call it a *single blow up*. A blow up is called *double⁺* blow up if the blown up point belongs to two components of $*(\omega_{Sh} + C)$ (this is the pushdown). Similarly define a *single⁺* blow up.

Lemma 1.6.10 For any exceptional $G_{\beta} \in \overline{exc}(\ddot{W}/Z)$ on a blow up model $\ddot{W}$ we have:

- ◇ $-1 + \delta \leq N_{\ddot{W},G_{\beta}}$ if $\overline{exc}(\ddot{W}/Z)_{G_{\beta}}$ is of type D_r, E_6, E_7 or E_8 . In particular, in these cases $-1 + \delta \leq N_{\ddot{W}}(G_{\beta})$ holds.
- ◇ $2(-1 + \delta) \leq N_{\ddot{W},G_{\beta}}$ and $-1 + \delta \leq N_{\ddot{W}}(G_{\beta})$ if $\overline{exc}(\ddot{W}/Z)_{G_{\beta}}$ is of type A_r except if it is strictly monotonic.

Proof D_r case: Similar to the notation in the proof of theorem 1.5.1 let $G_\beta, G_{\beta'}, G_{\beta_1}, \dots, G_{\beta_r}$ be the exceptional divisors in $\overline{ex\bar{c}}(\ddot{W}/Z)_{G_\beta}$. Then from the equations in the proof of theorem 1.5.1 for the D_r case we get the following system for the log discrepancies:

$$\begin{aligned}
2a_\beta - a_{\beta_1} - 1 &\leq 0 \\
2a_{\beta'} - a_{\beta_1} - 1 &\leq 0 \\
2a_{\beta_1} - a_\beta - a_{\beta'} - a_{\beta_2} + 1 &\leq 0 \\
2a_{\beta_2} - a_{\beta_1} - a_{\beta_3} &\leq 0 \\
&\vdots \\
2a_{\beta_{r-1}} - a_{\beta_{r-2}} - a_{\beta_r} &\leq 0 \\
2a_{\beta_r} - a_{\beta_{r-1}} - 1 &\leq 0
\end{aligned}$$

Adding the first and the second equations gives $2a_\beta + 2a_{\beta'} - 2a_{\beta_1} - 2 \leq 0$ and putting this in the third equation we get $a_{\beta_1} \leq a_{\beta_2}$ and so $a_{\beta_1} \leq a_{\beta_2} \leq \dots \leq a_{\beta_r}$. So

$$\begin{aligned}
N_{\ddot{W}, G_\beta} &\geq a_\beta + a_{\beta'} + a_{\beta_r} - a_{\beta_1} - 2 \geq a_\beta + a_{\beta'} + a_{\beta_2} - a_{\beta_1} - 2 \\
&\geq 2a_{\beta_1} + 1 - a_{\beta_1} - 2 \geq a_{\beta_1} - 1 \geq \delta - 1
\end{aligned}$$

because $2a_{\beta_1} + 1 \leq a_\beta + a_{\beta'} + a_{\beta_2}$ and the fact that X is δ -lc.

A_r case [non-strictly monotonic]: In this case assume that the exceptional divisors in $\overline{ex\bar{c}}(\ddot{W}/Z)_{G_\beta}$ are $G_{\beta_1}, \dots, G_{\beta_r}$ so we get the system:

$$\begin{aligned}
2a_{\beta_1} - a_{\beta_2} - 1 &\leq 0 \\
2a_{\beta_2} - a_{\beta_1} - a_{\beta_3} &\leq 0 \\
&\vdots \\
2a_{\beta_{r-1}} - a_{\beta_{r-2}} - a_{\beta_r} &\leq 0 \\
2a_{\beta_r} - a_{\beta_{r-1}} - 1 &\leq 0
\end{aligned}$$

So there will be k such that $a_{\beta_1} \geq a_{\beta_2} \geq \dots \geq a_{\beta_k} \leq a_{\beta_r}$. Thus $N_{\ddot{W}}(G_{\beta_1}) \geq a_{\beta_1} + a_{\beta_1} - a_{\beta_2} - 1 \geq a_{\beta_1} - 1 \geq \delta - 1$. In this way we get

the similar inequalities for all other equations except for $N_{\ddot{W}}(G_{\beta_k})$. Suppose $N_{\ddot{W}}(G_{\beta_k}) < \delta - 1$. So we get $2a_{\beta_k} - a_{\beta_{k-1}} - a_{\beta_{k+1}} < \delta - 1$ and so $1 - \delta < a_{\beta_{k-1}} + a_{\beta_{k+1}} - 2a_{\beta_k} \leq a_{\beta_1} + a_{\beta_r} - 2a_{\beta_k}$. On the other hand by adding all the equations in the system above we get $N_{\ddot{W}, G_\beta} \geq a_{\beta_1} + a_{\beta_r} - 2 > 1 - \delta + 2a_{\beta_k} - 2 \geq \delta - 1$. This is a contradiction with the fact that $N_{\ddot{W}}(G_{\beta_k}) \geq N_{\ddot{W}, G_{\beta_k}}$.

To get the inequality for $N_{\ddot{W}, G_{\beta_k}}$ add all the equations in the system above. Note that if $r = 2$ then $a_{\beta_1} = a_{\beta_2}$ and checking the lemma is easy in this case.

E_6, E_7, E_8 cases:^{1.6.1} In these cases the graph $\overline{exc}(\ddot{W}/Z)_{G_\beta}$ is as in the recall 1.5.7. It is enough to put 2 in place of all self-intersection numbers because the negativity becomes smaller. We start from the smallest possible graph i.e. Case 1 in the mentioned recall:

$$\begin{aligned} 2a_\beta - a_{\beta_2} - 1 &\leq 0 \\ 2a_{\beta_2} - a_\beta - a_{\beta_1} - a_{\beta_3} + 1 &\leq 0 \\ 2a_{\beta_1} - a_{\beta_2} - 1 &\leq 0 \\ 2a_{\beta_3} - a_{\beta_2} - 1 &\leq 0 \end{aligned}$$

Adding all the equations we get $N_{\ddot{W}, G_\beta} = a_\beta + a_{\beta_1} + a_{\beta_3} - a_{\beta_2} - 2$. And by the second equation we have $a_\beta + a_{\beta_1} + a_{\beta_3} - a_{\beta_2} \geq a_{\beta_2} + 1$ so $N_{\ddot{W}, G_\beta} \geq a_{\beta_2} + 1 - 2 \geq \delta - 1$. In order this was a special case of the D_r type (the similarity of the system not necessarily the graph $\overline{exc}(\ddot{W}/Z)_{G_\beta}$). Note that the inequality for the total negativity implies the inequality for the negativity of each exceptional curve.

Now we prove other cases by induction on the number of the exceptional curves. The minimum is 4 exceptional curves and we just proved this case. Suppose we have proved up to $k - 1$ and our graph has k members. Suppose the exceptional curves are $G_\beta, G_{\beta_1}, \dots, G_{\beta_{k-1}}$ and such that G_{β_l} cuts $G_\beta, G_{\beta_{l-1}}$ and $G_{\beta_{l+1}}$. If $l = 2$ or $l = k - 2$ then again this will be a system of type D_r . Assume that otherwise happens. Since $-a_\beta + 1 \geq 0$ so we get a system as follows

$$2a_{\beta_1} - a_{\beta_2} - 1 \leq 0$$

^{1.6.1}I just prove that $-1 + \delta \leq N_{\ddot{W}}(G)$ for any exceptional G . We won't need the inequality for total negativity.

$$2a_{\beta_2} - a_{\beta_3} - a_{\beta_1} \leq 0$$

$$\vdots$$

$$2a_{\beta_{k-1}} - a_{\beta_{k-2}} - 1 \leq 0$$

This is a system of type A_{k-1} so either we have $a_{\beta_1} \geq a_{\beta_2}$ or $a_{\beta_{k-1}} \geq a_{\beta_{k-2}}$. Assume that the first one holds (the other case is similar). Now note that $N_{\tilde{W}}(G_{\beta_1}) \geq 2a_{\beta_1} - a_{\beta_2} - 1 = a_{\beta_1} - a_{\beta_2} + a_{\beta_1} - 1 \geq \delta - 1$. By ignoring G_{β_1} we get a system for a graph with a smaller number of elements:

$$2a_{\beta_2} - a_{\beta_3} - 1 \leq 2a_{\beta_2} - a_{\beta_3} - a_{\beta_1} \leq 0$$

$$\vdots$$

$$2a_{\beta_l} - a_{\beta_{l-1}} - a_{\beta_{l+1}} - a_{\beta} + 1 \leq 0$$

$$\vdots$$

$$2a_{\beta_{k-1}} - a_{\beta_{k-2}} - 1 \leq 0$$

and the lemma is proved by induction. $\square$

Lemma 1.6.11 *Suppose $\xi \in \ddot{W}/\overline{W}$ ($\ddot{W}$ is a blow up model), $\tilde{W}$ the blow up of $\ddot{W}$ at ξ and G_α the exceptional divisor of the blow up. Then we have the followings:*

If G_α is the double blow up of G_β and G_γ (i.e. $\xi \in G_\beta \cap G_\gamma$) then:

- $\diamond$ $N_{\tilde{W}}(G_\alpha) = a_\alpha - a_\beta - a_\gamma$ where a_α is the log discrepancy of G_α for K_X and similarly a_β and a_γ .
- $\diamond$ $N_{\tilde{W}}(G_\beta) = N_{\ddot{W}}(G_\beta) - N_{\tilde{W}}(G_\alpha)$ and $N_{\tilde{W}}(G_\gamma) = N_{\ddot{W}}(G_\gamma) - N_{\tilde{W}}(G_\alpha)$.
- $\diamond$ $N_{\tilde{W}} = N_{\ddot{W}} - N_{\tilde{W}}(G_\alpha)$.

And if G_α is the single blow up of G_β then

- $\diamond$ $N_{\tilde{W}}(G_\beta) = N_{\ddot{W}}(G_\beta) - N_{\tilde{W}}(G_\alpha)$, $N_{\tilde{W}}(G_\alpha) = a_\alpha - a_\beta - 1 \leq -\delta$ and $N_{\ddot{W}}(G_\beta) + \delta \leq 0$.
- $\diamond$ $N_{\tilde{W}} = N_{\ddot{W}}$.

Proof Standard computations and left to the reader. $\square$

Corollary 1.6.12 *If G_α is a single blow up of G_β on $\ddot{W}$, a blow up model of $\overline{W}$, and $N_{\ddot{W}}(G_\beta) \geq \delta - 1$ then $a_\alpha \geq a_\beta + \delta$.*

Proof Since G_α is a single blow up of G_β then $1 + a_\beta - a_\alpha + N_{\ddot{W}}(G_\beta) \leq 0$ and so $1 + a_\beta - a_\alpha + \delta - 1 \leq 0$ then $a_\beta + \delta \leq a_\alpha$.

Definition 1.6.13 Let ξ be a point on a blow up model $\ddot{W}$. Define the *multiplicity of double blow ups* as

$$\mu_{db}(\xi) = \max\{\#\{\text{double blow ups } / \xi \text{ before having a single blow up } / \xi\}\}$$

the maximum is taken over all sequences of blow ups from $\ddot{W}$ to W . The next lemma shows the boundedness of this number.

Lemma 1.6.14 $\mu_{db}(\xi)$ is bounded.

Proof Since by lemma 1.6.11 each double blow up adds a non-negative number to the total negativity of the system and since the total negativity is bounded^{1.6.2} then except a bounded number of double blow ups we have $\frac{-\delta}{2} \leq N_{\ddot{W}}(G_\alpha) = a_\alpha - a_\beta - a_\gamma \leq 0$ where G_α is the double blow up of some G_β and G_γ and $G_\beta \cap G_\gamma / \xi$. The inequality shows that $a_\beta + \frac{\delta}{2} \leq a_\beta + a_\gamma - \frac{\delta}{2} \leq a_\alpha$ and similarly $a_\gamma + \frac{\delta}{2} \leq a_\alpha$. In other words the log discrepancy is increasing at least by $\frac{\delta}{2}$. Since log discrepancies are in $[\delta, 1]$ then the number of these double blow ups has to be bounded. $\square$

Definition 1.6.15 Let $\xi \in \ddot{W}$ a blow up model. Define the *single blow up multiplicity* of ξ as:

$$\mu_{sb}(\xi) = \max\{\#\{G : G \text{ is single blown up and } G/\xi\}\}$$

The maximum is taken over all sequences of blow ups from $\ddot{W}$ to W . In the above definition G is the exceptional divisor of a single blow up $/\xi$. Also define $\mu_{sb}(G_\beta) = \sum_{\xi \in G_\beta} \mu_{sb}(\xi)$ and $\mu_{sb}(\ddot{W}) = \sum_{\xi \in \ddot{W}} \mu_{sb}(\xi)$.

^{1.6.2}because the total negativity on $\overline{W}$ is bounded. This boundedness for the A_r and D_r cases is shown in lemma 1.6.10 and for other cases it is obvious

So if ξ_2/ξ_1 (these points may be on different models) then $\mu_{sb}(\xi_1) \geq \mu_{sb}(\xi_2)$.

Remark 1.6.16 Usually there is not a unique sequence of blow ups from $\ddot{W}$ to W . In order if $\xi_1 \neq \xi_2$ are points on $\ddot{W}$ and they are centres of some exceptional divisors on W then it doesn't matter which one we first blow up to get to W i.e. they are independent.

Definition 1.6.17 Let $\xi \in \overline{exc}(\ddot{W}/Z)$ be a point on a blow up model $\ddot{W}$. We call such a point a *base point* if there is an exceptional divisor G_α/ξ on a blow up model $\tilde{W}$ such that $N_{\tilde{W}}(G_\alpha) < \delta - 1$.

Remark 1.6.18 By lemma 1.6.10 and lemma 1.6.11 if $\xi \in \overline{exc}(\ddot{W}/Z)_{G_\beta}$ and $\overline{exc}(\ddot{W}/Z)_{G_\beta}$ is of type A_r (non-strictly monotonic) , D_r , E_6 , E_7 or E_8 then ξ can not be a base point. Moreover again by lemma 1.6.10 if $\overline{exc}(\ddot{W}/Z)_{G_\beta}$ is strictly monotonic then there can be at most one base point in $\overline{exc}(\ddot{W}/Z)_{G_\beta}$ and it only can belong to the base curve.

Lemma 1.6.19 $\mu_{sb}(\xi)$ is bounded if $\xi \in \ddot{W}$ is not a base point.

Proof If G_α/ξ is a single blown up exceptional divisor then since ξ is not a base point we get $\delta - 1 \leq N_{\tilde{W}}(G_\alpha)$. So if G_α is a single blow up/ ξ of G_β then $a_\alpha \geq a_\beta + \delta$ i.e. it increases the log discrepancy at least by δ . And as we said in the proof of lemma 1.6.14, except a bounded number, any other double blow up/ ξ increases the log discrepancy at least by $\frac{\delta}{2}$. In order, there can be only a bounded number of blow ups/ ξ from $\ddot{W}$ to W . $\square$

Corollary 1.6.20 The number of exceptional curves/ ξ on W is bounded for any non-base point $\xi \in \ddot{W}$.

We now continue to the proof of theorem 1.6.1. If no divisor in ω_{Sh} has coefficient 1 then this is what we are looking for. Since in this case $K_W + \omega_{Sh}$ will be a 1/6-lc log divisor. If the opposite happens i.e. some divisor appear with coefficient 1 in ω_{Sh} then these divisors will form a connected chain $\mathcal{C}$ which doesn't intersect any other exceptional divisor/ Z with positive coefficient in ω_{Sh} , except in the edges of this chain. Some of the exceptional divisors of type F may appear with positive coefficients and some with zero coefficients in ω_{Sh} .

The image of the graph $\mathcal{G}$ on $W_{\mathcal{G}}$, that is $\mathcal{G}_1$ (see definition 1.6.6), is either of type A_r , D_r , E_6 , E_7 or E_8 because similar to what we proved above for $\overline{W}$ the model $W_{\mathcal{G}}$ is the minimal resolution of some surface i.e. the minimal resolution of the surface $X_{\mathcal{G}}$ obtained from X by contracting the exceptional/ Z curves on X whose birational transform belong to $\mathcal{G}$. In order there is no -1 -curve/ $X_{\mathcal{G}}$ on $W_{\mathcal{G}}$.

Now suppose $\mathcal{G}_1$ is of type A_r , not strictly monotonic and let the pushdown of the chain $\mathcal{C}$ be $\mathcal{C}_1$ on $W_{\mathcal{G}}$. Let the exceptional divisors of $\mathcal{G}_1$ be $G_{\beta_1}, \dots, G_{\beta_r}$ and assume that the chain $\mathcal{C}_1$ consists of $G_{\beta_k}, \dots, G_{\beta_l}$. Hence $N_{W_{\mathcal{G}}}^{Sh}(G_{\beta_k}) \leq -\frac{1}{6}$, $N_{W_{\mathcal{G}}}^{Sh}(G_{\beta_{k+1}}) = \dots = N_{W_{\mathcal{G}}}^{Sh}(G_{\beta_{l-1}}) = 0$ and $N_{W_{\mathcal{G}}}^{Sh}(G_{\beta_l}) \leq -\frac{1}{6}$. Here the superscript Sh means that we compute the negativity according to the Shokurov log divisor not the primary log divisor (we have already defined it above). Note that if $a_{\beta}^{Sh} > 0$ for some β then $a_{\beta}^{Sh} \geq \frac{1}{6}$ because the denominator of a_{β}^{Sh} is in $\{1, 2, 3, 4, 6\}$. The chain $\mathcal{C}_1$ is of type A_{l-k+1} . From the constructions in the local isomorphism subsection we can replace the Shokurov log numbers $a_{\beta_k}^{Sh} = 0, \dots, a_{\beta_l}^{Sh} = 0$ with new log numbers with bounded denominators and preserve all other Shokurov log numbers in the graph $exc(W_{\mathcal{G}}/Z)$ such that we obtain a new log divisor $K_{W_{\mathcal{G}}} + \Omega_1$ on $W_{\mathcal{G}}$ which is antinef/ Z and is a Klt log divisor. Now put $K_W + \Omega =^* (K_{W_{\mathcal{G}}} + \Omega_1)$. The only problem with Ω is that it may have negative coefficients (it is a subboundary). Remark 1.6.18 and corollary 1.6.20 assure us that the negativity of these coefficients is bounded from below. Moreover if an exceptional divisor has negative coefficient in Ω then it should belong to the graph $\mathcal{G}$. But any exceptional divisor in $\mathcal{G}$ appears with positive coefficient in ω_{Sh} . Since $\omega_{Sh} \geq \omega$ and also by the definition of $\mathcal{G}$, any exceptional divisor of type F in $\mathcal{G}$ has positive coefficient at least $\frac{1}{6}$. And if E is not of type F but belongs to $\mathcal{G}$ then since B^+ is not zero $P \in Z$ we get positive coefficients in ω_{Sh} for all exceptional/ Z curves which are not of type F . Thus all members of $\mathcal{G} = Exc(W/Q)$ appear with positive coefficient in ω_{Sh} .

Now consider the sum

$$K_W + \Omega + I[K_W + \omega_{Sh}] = (1 + I)K_W + [\Omega + I\omega_{Sh}]$$

where I is an integer. Mentioning the fact that the negative coefficients appeared in Ω are bounded from below, this implies that there is a large bounded I such that the sum $\Omega + I\omega_{Sh}$ is an effective divisor. So by construction the log divisor $K_W + \frac{\Omega + I\omega_{Sh}}{1+I}$ is ϵ -lc and antinef/ Z for some fixed rational number $0 < \epsilon$ and the denominators of the coefficients in the log

divisor are bounded.

Now assume that $\mathcal{G}_1$ is strictly monotonic and the base curve is G_{β_1} . By corollary 1.6.20 and remark 1.6.18 the only place where we may have difficulties is the base point, ξ , on the base curve if there is any such point.

Now we blow ξ up and get the exceptional divisor G_{α_1} . The chain $G_{\alpha_1}, G_{\beta_1}, \dots, G_{\beta_r}$ is not exactly of type A_{r+1} because G_{α_1} is a -1 -curve. But still we can claim that there is at most a base on this chain and it only can be on G_{α_1} . Obviously a base point cannot be on $G_{\beta_2}, \dots, G_{\beta_r}$. Now suppose that the intersection point of G_{α_1} and G_{β_r} is a base point. Then the sum of negativities of all $G_{\alpha_1}, G_{\beta_1}, \dots, G_{\beta_r}$ should be less than $2\delta - 2$. This is impossible because the sum of negativities of all $G_{\beta_1}, \dots, G_{\beta_r}$ on $W_{\mathcal{G}}$ is at least $2\delta - 2$ (remember that blowing up reduces negativity).

Now if on G_{α_1} there is a base point ξ_1 then again we blow this point up to get G_{α_2} and so on. This process has to stop after finitely many steps (not after bounded steps!). Let the final model to be W_{ξ} and $G_{\alpha_1}, \dots, G_{\alpha_s}$ the new exceptional divisors. In order we have constructed a chain (because on each curve there was at most one base point) and by adding the new exceptional divisors to $\mathcal{G}_1$ we get a new graph $\mathcal{G}_2$. Now there is no base point on $\mathcal{G}_2$. All the divisors G_{α_i} have self-intersection equal to -2 except G_{α_s} which is a -1 -curve.

Now let $\mathcal{C}_2$ to be the pushdown of $\mathcal{C}$ i.e. the connected chain of curves with coefficient one in ω_{Sh} on W_{xi} . If G_{α_s} is not in $\mathcal{C}_2$ then we proceed exactly as in the non-monotonic case above; that is we assign appropriate coefficients to the members of $\mathcal{C}_2$ and keep all other coefficients in ω_{Sh} on W_{xi} . If G_{α_s} is in $\mathcal{C}_2$ then let $\mathcal{C}'$ be the chain $\mathcal{C}_2$ except the member G_{α_s} . This new chain (i.e. $\mathcal{C}'$) is of type A_x and so we can assign appropriate coefficients to its members and put the coefficient of G_{α_s} simply equal to zero and keep all other coefficients in ω_{Sh} on W_{xi} . In any case we construct a Klt log divisor $K + \Omega$ on W_{xi} which is antinef/ Z and the boundary coefficients are with bounded denominators. The rest is as in the non-monotonic case above.

Suppose the graph $\mathcal{G}_1$ is of type D_r and $\mathcal{C}_1 \neq \emptyset$ (if it is empty then we already have Ω_1). Assume that the members of $\mathcal{G}_1$ are $G_{\beta}, G_{\beta'}, G_{\beta_1}, \dots, G_{\beta_r}$ and the members of $\mathcal{C}_1$ are $G_{\beta_k}, \dots, G_{\beta_l}$. As in the proof of lemma 1.6.10 for the D_r case we have $a_{\beta_1}^{Sh} \leq a_{\beta_2}^{Sh} \leq \dots$. So $k = 1$ and we have $2a_{\beta}^{Sh} - 0 - 1 \leq 0$ and so $a_{\beta}^{Sh} \leq \frac{1}{2}$ and similarly $a_{\beta'}^{Sh} \leq \frac{1}{2}$. The chain $\mathcal{C}_1$ is of type A_l and so we can change the coefficients of its members in ω_{Sh} on $W_{\mathcal{G}}$. The rest of the argument is very similar to the above cases. Just note that there is no base

point in this case.

The cases E_6 , E_7 and E_8 are safe by remark 1.6.18 and corollary 1.6.20. In these cases the graph $\mathcal{G}$ is bounded so assigning the primary log numbers to the members of $\mathcal{G}_1$ and Shokurov log numbers to the rest of the graph $\text{exc}(W_{\mathcal{G}}/Z)$ gives a log divisor which can be used as $K_{W_{\mathcal{G}}} + \Omega_1$. Here the proof of theorem 1.6.1 is finished.

□

1.7 Global case

The main theorem of this subsection is the following theorem. A generalised version of this and the BAB follow as corollaries.

Theorem 1.7.1 *Conjecture $WC_{\delta,2,\{0\}}$ holds in the global case i.e. when Z is a point.*

Proof We divide the problem into two main cases: exceptional and non-exceptional. $(X, 0)$ is non-exceptional if there is a non-Klt $\mathbb{Q}$ -complement $K_X + M$. By [Sh2, 2.3.1], under our assumptions on X , non-exceptionality is equivalent to the fact that K_X has a non-Klt $(0, n)$ -complement for some $n < 58$. We prove that the exceptional cases are bounded. But in the non-exceptional case we only prove the existence of an (ϵ, n) -complement for a bounded n . Later we show that this in order implies the boundedness of X .

First assume that $(X, 0)$ is **non-exceptional**.

1. Lets show the set of accumulation points of the mlds in dim 2 for lc pairs (T, B) where $B \in \Phi_{sm}$, by $\text{Accum}_{2, \Phi_{sm}}$. Then $\text{Accum}_{2, \Phi_{sm}} \cap [0, 1] = \{1 - z\}_{z \in \Phi_{sm}} = \{\frac{1}{k}\}_{k \in \mathbb{N}} \cup \{0\}$ [Sh8]. Now if there is a $\tau > 0$ such that $\text{mld}(P, T, B) \notin [\frac{1}{k}, \frac{1}{k} + \tau]$ for any natural number k and any point $P \in T$ then there will be only a finite number of possibilities for the index of $K_T + B$ at P if (T, B) is $\frac{1}{m}$ -lc for some $m \in \mathbb{N}$. Now Borisov-Mckernan [Mc, 1.2] implies the boundedness of all such T if $-(K_T + B)$ is nef and big and τ and m are fixed. In order in the following steps we try to reduce our problem to this situation in some cases.

2. **Definition 1.7.2** Let $B = \sum b_i B_i$ be a boundary on a variety T and $\tau > 0$ a real number. Define

$$D_\tau := \sum_{b_i \notin [\frac{k-1}{k} - \tau, \frac{k-1}{k}]} b_i B_i + \sum_{b_i \in [\frac{k-1}{k} - \tau, \frac{k-1}{k}]} \frac{k-1}{k} B_i$$

where in the first term $b_i \notin [\frac{k-1}{k} - \tau, \frac{k-1}{k}]$ for any natural number k but in the second term k is the smallest natural number satisfying $b_i \in [\frac{k-1}{k} - \tau, \frac{k-1}{k}]$.

Lemma 1.7.3 For any natural number m there is a real number $\tau > 0$ such that if (T, B) is a surface log pair, $P \in T$, $K_T + B$ is $\frac{1}{m}$ -lc at P and $D_\tau \in \Phi_{sm}$ then $K_T + D_\tau$ is also $\frac{1}{m}$ -lc at P .

Note that τ depends only on m .

Proof By applying the ACC to all surface pairs with standard boundary, we get a fixed rational number $v > 0$ such that if any $K_T + D_\tau$ is not $\frac{1}{m}$ -lc at P then $\text{mld}(P, T, D_\tau) < \frac{1}{m} - v$.

Now assume that the lemma is not true. So there is a sequence $\tau_1 > \tau_2 > \dots$ and a sequence of pairs $\{(T_i, B_i)\}$ where if we take τ_i for the pair (T_i, B_i) then the lemma doesn't hold at $P_i \in T_i$. In other words $\text{mld}(P_i, T_i, D_{\tau_i}) < \frac{1}{m} - v$.

Write $B_i := F_i + C_i$ where $F_i = \sum f_{i,x} F_{i,x}$ and $C_i = \sum c_{i,y} C_{i,y}$ have no common components and the coefficient of any component of C_i is equal to the coefficient of the same component in D_{τ_i} but the coefficient of any component of F_i is less than the coefficient of the same component in D_{τ_i} .

Now there is a set $\{s_{1,x}\} \subseteq [\frac{m-1}{m} - \tau_1, \frac{m-1}{m}]$ of rational numbers such that $\text{mld}(P_1, T_1, \sum s_{1,x} F_{1,x} + C_1) = \frac{1}{m} - v$. There is i_2 such that $\max\{s_{1,x}\} < \frac{m-1}{m} - \tau_{i_2}$. So there is also a set $\{s_{2,x}\} \subseteq [\frac{m-1}{m} - \tau_{i_2}, \frac{m-1}{m}]$ such that $\text{mld}(P_{i_2}, T_{i_2}, \sum s_{2,x} F_{i_2,x} + C_{i_2}) = \frac{1}{m} - \frac{v}{2}$. By continuing this process we find $\{s_{j,x}\} \subseteq [\frac{m-1}{m} - \tau_{i_j}, \frac{m-1}{m}]$ such that $\max\{s_{i_{j-1},x}\} < \frac{m-1}{m} - \tau_{i_j}$. Hence we can find a set $\{s_{j,x}\} \subseteq [\frac{m-1}{m} - \tau_{i_j}, \frac{m-1}{m}]$ such that $\text{mld}(P_{i_j}, T_{i_j}, \sum s_{j,x} F_{i_j,x} + C_{i_j}) = \frac{1}{m} - \frac{v}{j}$.

In order we have constructed a set $\cup\{s_{j,x}\}$ of rational numbers which satisfies the DCC condition but there is an increasing set of mlds corresponding to boundaries with coefficients in $\cup\{s_{j,x}\}$. This is a contradiction with the ACC for mlds. $\square$

3. Let m be the smallest number such that $\frac{1}{m} \leq \delta$. Let $h = \min\{\frac{k-1}{k} - \frac{u}{r!} > 0\}_{1 \leq k \leq m}$ where u, k are natural numbers and $r = \max\{m, 57\}$. Now choose a τ for m as in lemma 1.7.3 such that $\tau < h$.

Blow up one exceptional divisor E via $f : Y \longrightarrow X$ such that the log discrepancy satisfies $\frac{1}{k} \leq a(E, X, 0) \leq \frac{1}{k} + \tau$ for some $k > 1$ (if such E doesn't exist then go to step 1). The crepant log divisor $K_Y + B_Y$ is $\frac{1}{m}$ -lc and so by lemma 1.7.3 $K_Y + D_\tau$ is also $\frac{1}{m}$ -lc (D_τ is constructed for B_Y). Let $K_X + B^+$ be a $(0, n)$ -complement for some $n < 58$ and $K_Y + B_Y^+$ be the crepant blow up. Then by the way we chose τ we have $D_\tau \leq B^+$. Now run the anti-LMMP over $K_Y + D_\tau$ i.e. contract any birational type extremal ray R such that $(K_Y + D_\tau).R > 0$. At the end of this process we get a model X_1 and the corresponding map $g : Y \longrightarrow X_1$. After contracting those birational extremal rays where $K_{X_1} + D_\tau$ is numerically zero we get a model S_1 with one of the following properties:

- ◇ $\rho(S_1) = 1$ and $K_{S_1} + D_\tau \equiv K_{S_1} + B_{S_1}^+ \equiv 0$ and $\frac{1}{m}$ -lc.
- ◇ $\rho(S_1) = 2$ and $(K_{S_1} + D_\tau).R = 0$ for a non-birational type extremal ray R on S_1 and $K_{S_1} + D_\tau$ is $\frac{1}{m}$ -lc.
- ◇ $-(K_{S_1} + D_\tau)$ is nef and big and $K_{S_1} + D_\tau$ is $\frac{1}{m}$ -lc.

where $K_{S_1} + D_\tau$ is the birational transform of $K_Y + D_\tau$.

In any case $-(K_{S_1} + D_\tau)$ is nef because $D_\tau \leq B_{S_1}^+$ and so D_τ can not be positive on a non-birational extremal ray. $K_{S_1} + D_\tau$ is $\frac{1}{m}$ -lc by the way we have chosen τ .

4. If the first case occurs in the division in step 3 then we are done.
5. If the second case occurs in the division in step 3 then R defines a fibration $\phi : S_1 \longrightarrow Z$. Note that $B_{S_1}^+ = D_\tau + N$ where each component of N is a fibre of ϕ and there are only a finite number of possibilities for the coefficients of N . Now we can replace N by $N' \equiv N$ where each

component of N' is a general fibre of ϕ , with only a finite number of possibilities for the coefficients of N' and such that $K_{S_1} + D_\tau + N'$ is $\frac{1}{m}$ -lc. Note that the components of N' are smooth curves and intersect the components of D_τ transversally in smooth points of S_1 . Now the only problem is that we don't know if the index of $K_{S_1} + D_\tau + N'$ is bounded or not. Note that it is enough if we can get the boundedness of the index of $K_{S_1} + D_\tau$.

6. Now assume that the third case or the second case occurs in the division in step 3. Let C be a curve contracted by $g : Y \rightarrow X_1$ constructed in step 3. If C is not a component of B_Y then the log discrepancy of C with respect to $K_{X_1} + B_{X_1}$ is at least 1 where $K_{X_1} + B_{X_1}$ is the birational transform of $K_Y + B_Y$. Moreover $g(C) \in \text{Supp } B_{X_1} \neq \emptyset$. So the log discrepancy of C with respect to K_{X_1} is more than 1. This means that C is not a divisor on a minimal resolution $W_1 \rightarrow X_1$. Let $W \rightarrow X$ be a minimal resolution. Then there is a morphism $W \rightarrow W_1$. Hence $\text{exc}(W_1/X_1) \subseteq \text{exc}(W/X)$. Now if $C \in \text{exc}(W/X)$ is exceptional/ X_1 then $a(C, X_1, D_\tau) < a(C, X, 0)$.
7. Let $(X_1, B_1) := (X_1, D_\tau)$ and repeat the process. In other words again we blow up one exceptional divisor E via $f_1 : Y_1 \rightarrow X_1$ such that the log discrepancy satisfies $\frac{1}{k} \leq a(E, X_1, B_1) \leq \frac{1}{k} + \tau$ for some natural number $k > 1$. The crepant log divisor $K_{Y_1} + B_{1,Y_1}$ is $\frac{1}{m}$ -lc and so by lemma 1.7.3 $K_{Y_1} + D_{1,\tau}$ is $\frac{1}{m}$ -lc. Note that the point which is blown up on X_1 can not be smooth since $\tau < h$ as defined in step 3. So according to step 6 the blown up divisor E is a member of $\text{exc}(W/X)$. Now we again run the anti-LMMP on $K_{Y_1} + D_{1,\tau}$ and proceed as in step 3.

$$\begin{array}{ccccccc}
W & \longrightarrow & W_1 & \longrightarrow & W_2 & \longrightarrow & \cdots \\
\downarrow & & \downarrow & & \downarrow & & \\
Y & & Y_1 & & Y_2 & & \cdots \\
\downarrow f & \searrow g & \downarrow f_1 & \searrow g_1 & \downarrow & \searrow & \\
X & & X_1 & & X_2 & & \cdots \\
& & \downarrow & & \downarrow & & \\
& & S_1 & & S_2 & & \cdots
\end{array}$$

8. Steps 6,7 show that each time we blow up a member of $\text{exc}(W/X)$ say E . And if we blow that divisor down in some step then the log discrepancy $a(E, X_j, B_j)$ will decrease. That divisor will not be blown up again unless the log discrepancy drops at least by $\frac{1}{2(m-1)} - \frac{1}{2m}$. So after finitely many steps either the case one occurs in the division in step 3 or we get a model X_i with a standard boundary B_i such that there is no E where $\frac{1}{k} \leq a(E, X_i, B_i) \leq \frac{1}{k} + \tau$ for any $1 < k \leq m$. This later implies the boundedness of the index of $K_{X_i} + B_i = K_{X_i} + D_{i-1, \tau}$. If $-(K_{X_i} + B_i)$ is nef and big (case one) then (X_i, B_i) will be bounded by step 1. Otherwise we have the second case in the division above and so by step 5 we are done (the index of $K_{X_i} + D_{i-1, \tau} + N'$ is bounded).

Now we treat the **exceptional** case: From now on we assume that $(X, 0)$ is exceptional.

9. Let $W \rightarrow X$ be a minimal resolution. Let $0 < \tau < \frac{1}{2}$ be a number and the minimal log discrepancy of $(X, 0)$ be $a = \text{mld}(X, 0)$. If $a \geq \frac{1}{2} + \tau$ then we know that X belongs to a bounded family according to step 1 above. So we assume $a < \frac{1}{2} + \tau$ and then blow up an exceptional/ X curve E_1 with log discrepancy $a_{E_1} = a(E_1, X, 0) \leq \frac{1}{2} + \tau$ to get $Y \rightarrow X$ and put $K_Y + B_Y = {}^*K_X$. Let $t \geq 0$ be a number such that there is an extremal ray R such that $(K_Y + B_Y + tE_1).R = 0$ and $E_1.R > 0$ (and s.t. $K_Y + B_Y + tE_1$ Klt and antinef). Such R exists otherwise there is a $t > 0$ such that $K_Y + B_Y + tE_1$ is lc (and not Klt) and antinef. This is a contradiction by [Sh2, 2.3.1]. Now contract $R : Y \rightarrow Y_1$ if it is of birational type.

Again by increasing t there will be an extremal ray R_1 on Y_1 such that $(K_{Y_1} + B_{Y_1} + tE_1).R_1 = 0$ and $E_1.R_1 > 0$ (preserving the nefness of $-(K_{Y_1} + B_{Y_1} + tE_1)$). If it is of birational then contract it and so on. After finitely many steps we get a model $(V_1, B_{V_1} + t_1E_1)$ and a number $t_1 > 0$ with the following possible outcomes:

(1.7.3.1)

- ◇ $(V_1, B_{V_1} + t_1E_1)$ is Klt, $\rho(V_1) = 1$ and $K_{V_1} + B_{V_1} + t_1E_1$ is antinef.
- ◇ $(V_1, B_{V_1} + t_1E_1)$ is Klt and $\rho(V_1) = 2$ and there is a non-birational extremal ray R on V_1 . Moreover $K_{V_1} + B_{V_1} + t_1E_1$ and K_{V_1} are antinef.

- ◇ $(V_1, B_{V_1} + t_1 E_1)$ is Klt and $\rho(V_1) = 2$ and there is a non-birational extremal ray R on V_1 . Moreover $K_{V_1} + B_{V_1} + t_1 E_1$ is antinef but K_{V_1} is not antinef.

Define $K_{V_1} + D_1 = K_{V_1} + B_{V_1} + t_1 E_1$. Note that in all the cases above E_1 is a divisor on V_1 and the coefficients of B_{V_1} and D_1 are $\geq \frac{1}{2} - \tau$.

Lemma 1.7.4 *Let $P \in U$ be a δ -lc surface singularity. Moreover suppose that there is at most one exceptional/ U divisor such that $a(E, U, 0) < \frac{1}{2} + \tau$. Then the index K_U is bounded at P where the bound only depends on δ and τ .*

Proof We only need to prove this when the singularity is of type A_r (otherwise the index is bounded). If there is no E/P such that $a(E, U, 0) < \frac{1}{2} + \frac{\tau}{2}$ then step 1 shows that the index is bounded. But if there is one E/P such that $a(E, U, 0) < \frac{1}{2} + \frac{\tau}{2}$ then using the notation as in 1.5.2.1 we have $a_{i+1} - a_i \geq \frac{\tau}{2}$ and $a_{i-1} - a_i \geq \frac{\tau}{2}$. This implies the boundedness of r and so the index at P . $\square$

10. Let U be a surface with the following properties:

- ◇ $\rho(U) = 1$.
- ◇ $K_U + G_U$ antinef, Klt and exceptional.
- ◇ K_U antiample.

Now blow up two divisors E and E' as $f : Y_U \rightarrow U$ such that $a(E, U, 0) < \frac{1}{2} + \tau$ and $a(E', U, 0) < \frac{1}{2} + \tau$ (suppose there are such divisors). Choose $t, t' \geq 0$ such that $(f^*(K_U + G_U) + tE + t'E') \cdot R = 0$ for an extremal ray R s.t. $R \cdot E \geq 0$ and $R \cdot E' \geq 0$ and $f^*(K_U + G_U) + tE + t'E'$ is antinef and Klt. We contract R to get $g : Y_U \rightarrow U'$. We call such operation a **hat of first type**. Note that E and E' are divisors on U' and $\rho(U') = 2$. Define $K_{U'} + G_{U'}$ to be the pushdown of $f^*(K_U + G_U) + tE + t'E'$.

If K_U is δ -lc and such E, E' don't exist as above then the index of K_U will be bounded by lemma 1.7.4. So U will be bounded.

11. Let U be a surface with the following properties:

- ◇ $\rho(U) = 2$.
- ◇ $K_U + G_U$ antinef, Klt and exceptional.
- ◇ $-K_U$ nef and big.

Now blow up a divisor E to get $f : Y_U \longrightarrow U$ such that $a(E, U, 0) < \frac{1}{2} + \tau$ (suppose there is such E). Let $t \geq 0$ be such that $(f^*(K_U + G_U) + tE).R = 0$ for an extremal ray R s.t. $R.E \geq 0$ and $f^*(K_U + G_U) + tE$ is antinef and Klt. We contract R to get $g : Y_U \longrightarrow U'$. We call such operation a **hat of second type**. Note that E is a divisor on U' and $\rho(U') = 2$. Define $K_{U'} + G_{U'}$ to be the pushdown of $f^*(K_U + G_U) + tE$. If K_U is δ -lc and such E doesn't exist as above then the index of K_U and so U will be bounded by lemma 1.7.4.

12. Let U be a surface with the following properties:

- ◇ $\rho(U) = 2$ and U is Pseudo-WLF.
- ◇ There is a birational type extremal ray R_{bir} and the other extremal ray of U is of fibration type.
- ◇ $K_U + G_U$ antinef, Klt and exceptional.
- ◇ $K_U.R_{bir} > 0$.

Then we say that U is of **2-bir** type. Let C be the divisor that defines R_{bir} on U . There is a $c \in (0, 1)$ such that $(K_U + cC).C = 0$. Now blow up E as $Y_U \longrightarrow U$ such that $a(E, U, cC) < \frac{1}{2} + \tau$ (suppose there is such E). Now let $t \geq 0$ such that $f^*(K_U + G_U + tC).R = 0$ for an extremal ray R s.t. $R.E \geq 0$, $R.C \geq 0$ and $f^*(K_U + G_U + tC)$ is antinef and Klt. We contract R to get $g : Y_U \longrightarrow U'$. We call such operation a **hat of third type**. Define $K_{U'} + G_{U'}$ to be the pushdown of $f^*(K_U + G_U + tC)$. Note that in this case E and C are both divisors on U' and $\rho(U') = 2$.

If $K_U + cC$ is δ -lc and such E doesn't exist as above then contract $C : U \longrightarrow U_1$. Thus the index of K_{U_1} will be bounded at each point by lemma 1.7.4 and so U_1 and consequently U will be bounded.

$$\begin{array}{ccc}
 Y_U & & \\
 \downarrow f & \searrow g & \\
 U & & U'
 \end{array}$$

13. Let U be a surface such that $\rho(U) = 2$ and $K_U + G_U$ antinef, Klt and exceptional where $G_U \neq 0$. Moreover suppose there are two exceptional curves H_1 and H_2 on U . In this case let C be a component of G_U and let $t \geq 0$ such that $(K_U + G_U + tC).H_i = 0$ for $i = 1$ or 2 and $K_U + G_U + tC$ Klt and antinef (assume $i = 1$). We contract H_1 as $U \longrightarrow U_1$ and define $K_{U_1} + G_{U_1}$ to be the pushdown of $K_U + G_U + tC$.

Definition 1.7.5 Define $K_U + \Delta_U$ as follows: $K_U + \Delta_U := K_U$ in step 10 and step 11. $K_U + \Delta_U := K_U + cC$ in step 12. And $K_{U_1} + \Delta_{U_1} := K_{U_1}$ in step 13.

14. The following lemmas are crucial to our proof.

Lemma 1.7.6 *Let $\mathcal{U}$ be a bounded family of surfaces with Picard number one or two and let $0 < x < 1$ be a rational number. Moreover assume the following for each member U_i :*

- ◇ $-(K_{U_i} + B_i)$ is nef and big for a boundary B_i where each coefficient of B_i is $\geq x$.
- ◇ $K_{U_i} + B_i$ is Klt.

Then $(U_i, \text{Supp } B_i)$ is bounded.

Proof In order we prove that there is a finite set Λ_f such that for each U_i there is a boundary $M_i \in \Lambda_f$ s.t. $-(K_{U_i} + M_i)$ is nef and big and $M_i \leq B_i$.

If $\rho(U_i) = 1$ then simply take $M_i = x \sum_{\alpha} B_{\alpha}$ where $B_i = \sum_{\alpha} B_{\alpha}$. Obviously $-(K_{U_i} + M_i)$ is nef and big and since U_i belongs to a bounded family so $(U_i, \text{Supp } M_i)$ is also bounded.

Now suppose $\rho(U_i) = 2$. Put $N_i = x \sum_{\alpha} B_{\alpha}$. If $-(K_{U_i} + N_i)$ is not nef then there should be an exceptional curve E on U_i where $(K_{U_i} + N_i).E > 0$. Let $\theta : U_i \longrightarrow U'_i$ be the contraction of E . By our assumptions $K_{U'_i} + B'_i$, the pushdown of $K_{U_i} + B_i$, is antiample. So $K_{U'_i} + N'_i$, the pushdown of $K_{U_i} + N_i$ is also antiample. Boundedness of U_i implies the boundedness of U'_i (since we have a bound for the Picard number of a minimal resolution of U'_i). Thus $-(K_{U_i} + M_i) := -\theta^*(K_{U'_i} + N'_i) = -(K_{U_i} + N_i + yE)$ is nef and big and there are only a finite number of

possibilities for $y > 0$. This proves the boundedness of $(U_i, \text{Supp}(N_i + yE))$. Note that in the arguments above $\text{Supp } B_i = \text{Supp } M_i$.

□

Lemma 1.7.7 (The main lemma) *Suppose that $\mathcal{U} = \{(U, \text{Supp } D)\}$ is a bounded family of log pairs of dim d where $K_U + D$ is antinef and ϵ -lc for a fixed $\epsilon > 0$. Then the set of partial resolutions of all $(U, D) \in \mathcal{U}$ is a bounded family.*

Note that here we don't assume (U, D) to be bounded i.e. the coefficients of D may not be in a finite set.

Proof Let (U_t, D_t) be a member of the family. By our assumptions the number of components of D_t is bounded (independent of t) and so we can consider any divisor supported in D_t as a point in a real finite dimensional space. Let $D_t = \sum_{1 \leq i \leq q} d_{i,t} D_{i,t}$ and define

$$\mathcal{H}_t := \{(h_1, \dots, h_q) \in \mathbb{R}^q \mid K_{U_t} + \sum_{1 \leq i \leq q} h_i D_{i,t} \text{ is antinef and } \epsilon - \text{lc}\}$$

So $\mathcal{H}_t$ is a subset of the cube $[0, 1]^q$ and since being ϵ -lc and antinef are closed conditions then $\mathcal{H}_t$ is a closed and hence compact subset of $[0, 1]^q$. In order $\{(U_t, \mathcal{H}_t)\}$ is a bounded family. For each $H \in \mathcal{H}_t$ the corresponding pair (U_t, H) is ϵ -lc. Let $Y_H \rightarrow U_t$ be a terminal blow up of (U_t, H) and assume that the set of exceptional/ U_t divisors on Y_H is R_H . For different H we may have different R_H but the union of all R_H is a finite set where H runs through $\mathcal{H}_t$. Suppose otherwise so there is a sequence $\{H_1, \dots, H_m, \dots\} \subseteq \mathcal{H}_t$ such that the union of all R_{H_i} is not finite. Since $\mathcal{H}_t$ is compact then there is at least an accumulation point in $\mathcal{H}_t$, say $\bar{H}$, for the sequence (we can assume that this is the only accumulation point). So $(U_t, \bar{H})$ is ϵ -lc. Let $v = (1, \dots, 1) \in \mathbb{R}^q$. Then there are $\alpha, \beta > 0$ such that $K_{U_t} + H_\alpha$ is $\epsilon - \beta$ -lc where $\epsilon - \beta > 0$ and H_α is the corresponding divisor of $\bar{H} + \alpha v$. In particular this implies that there is a (with positive radius) d -dim disc $\mathbb{B} \subseteq [0, 1]^q$ with $\bar{H}$ as its centre such that $K_{U_t} + H$ is $\epsilon - \beta$ -lc and $R_H \subseteq R_{H_\alpha}$ for any $H \in \mathbb{B}$. This is a contradiction with the way we chose the

sequence $\{H_1, \dots, H_m, \dots\}$. The function $R : \mathcal{H}_t \longrightarrow \mathbb{N}$ gives a finite decomposition of the set $\mathcal{H}_t$. This means that there are only a finite number of partial resolutions for all (U_t, H) where $H \in \mathcal{H}_t$ for a fixed t . Using Noetherian induction completes the proof. $\square$

Now we prove a statement similar to [Sh2, 4.2].

Lemma 1.7.8 *Let $\mathcal{U} = \{(U, \text{Supp } D)\}$ be a bounded family where we assume that each (U, D) is Klt and exceptional and $K + D$ is antinef. Then the singularity is bounded i.e. there is a constant $\gamma > 0$ such that each (U, D) is γ -lc.*

Proof For $(U, \text{Supp } B_i)$ a member of the family let

$$\mathcal{H}_i = \{H = \sum h_{k,i} D_{k,i} \mid K + H \text{ is log canonical and } -(K + H) \text{ is nef}\}$$

where $D_i = \sum d_{k,i} D_{k,i}$.

It is a closed subset of a multi-dimensional cube (with bounded dimension) and so it is compact. Let $a_i = \inf\{\text{mld}(U_i, H) : H \in \mathcal{H}_i\} > 0$. Since the family is bounded then $\{a_i\}$ is bounded from below.

$\square$

Now we return to the division in 1.7.3.1 and deal with each case as follows:

15. (First case in 1.7.3.1) Perform a hat of the first type for $U := V_1$ and $K_U + G_U := K_{V_1} + D_1$ (so we blow up E, E'). Then we get $V_2 := U'$ and $K_{V_2} + D_2 := K_{U'} + G_{U'}$ as defined above and $Y_1 := Y_U$. Now V_2 would be as in step 11, 12 or 13 so we can perform the appropriate operation as explained in each case. If V_2 is as in step 11 then $a(E, V_2, \Delta_{V_2}) = 1$ and $a(E', V_2, \Delta_{V_2}) = 1$. If V_2 is as in step 12 then E or E' is not exceptional so we have $a(E, V_2, \Delta_{V_2}) = 1$ or $a(E', V_2, \Delta_{V_2}) = 1$. But if V_2 is as in step 13 then we get U_1 as defined in step 13 and so $a(E, U_1, \Delta_{U_1}) = 1$ or $a(E', U_1, \Delta_{U_1}) = 1$. In the later case we define (replace) $(V_2, D_2) := (U_1, G_{U_1})$.

So whatever case we have for V_2 we have $a(A, V_2, \Delta_{V_2}) = 1$ at least for one $A \in \text{exc}(Y/X)$.


16. (Second case in 1.7.3.1) Here we perform a hat of second type for $U := V_1$ and $K_U + G_U := K_{V_1} + D_1$ to get $V_2 := U'$ and $K_{V_2} + D_2 := K_{U'} + G_{U'}$. If V_2 is as in step 11 then $a(E, V_2, \Delta_{V_2}) = 1$. If V_2 is as in step 12 then go to step 17. But if V_2 is as in step 13 then we get U_1 as defined in step 13 where $K_U + G_U := K_{V_2} + D_2$ and then continue the process for U_1 as in step 15.

Here in some cases we may not be able to make the singularities better for $K + \Delta$ immediately on V_2 but the algorithm ensures us that we will be able to do that in later steps.

17. (Third case in 1.7.3.1) In this case V_1 is 2-bir. We perform a hat of the third type where $U := V_1$ and $K_U + G_U := K_{V_1} + D_1$ so we get $V_2 := U'$ and $Y_1 := Y_U$ and $K_{V_2} + D_2 := K_{U'} + G_{U'}$. If V_2 is as in step 11 then $a(E, V_2, \Delta_{V_2}) = 1$ and $a(C, V_2, \Delta_{V_2}) = 1$ (E is the blown divisor and C is on V_1 , as in step 12 for $U := V_1$). If V_2 is as in step 12 then $a(E, V_2, \Delta_{V_2}) = 1$ or $a(C, V_2, \Delta_{V_2}) = 1$. Now if V_2 is as in step 13 then we get U_1 as defined in step 13 and so $a(E, U_1, \Delta_{U_1}) = 1$ or $a(C, U_1, \Delta_{U_1}) = 1$. Then we define (replace) $(V_2, D_2) := (U_1, G_{U_1})$.

So whatever case we have for V_2 we have $a(A, V_2, \Delta_2) = 1$, after the appropriate operations, at least for one $A \in \text{exc}(Y/X)$.

18. After finitely many steps we get V_r where W/V_r such that $K_W + D := *(K_{V_r} + D_r)$ with effective D where V_r is bounded. Since all the coefficients of B_{V_r} are $\geq \frac{1}{2} - \tau$ (B_{V_r} is the birational transform of B_W where $K_W + B_W = *K_X$) then (V_r, B_{V_r}) is also bounded by lemma 1.7.6. By construction $\text{Supp } D_r = \text{Supp } B_{V_r}$ and so (V_r, D_r) is bounded. Lemma 1.7.7 implies the boundedness of W and so of X .



□

Corollary 1.7.9 *The BAB Conjecture(1.1.4) holds in dim 2.*

Proof Reduction to the case $B = 0$: We run the anti-LMMP on the divisor K_X ; if there is an extremal ray R such that $K_X.R > 0$ then contract R to get $X \rightarrow X_1$. Note that $B.R < 0$ so the bigness of $-K_X$ will be preserved (So R has to be of birational type). Repeat the same process for X_1 i.e. if there is an extremal ray R_1 such that $K_{X_1}.R_1 > 0$ then contract it and so on. Since in each step we get a Pseudo-WLF then the canonical class can not become nef. Let $\overline{X}$ be the last model in our process, then $-K_{\overline{X}}$ is nef ad big. Now the boundedness of $\overline{X}$ implies the boundedness of X . So we replace X by $\overline{X}$ i.e. from now on we can assume $B = 0$.

By theorem 1.7.1 $(X, 0)$ has an (ϵ, n) -complement $K_X + B^+$ for some $n \in \mathcal{N}_\delta, 2, \{0\}$. Now let $W \rightarrow X$ be a minimal resolution and $\phi : W \rightarrow S$ be the map obtained by running the classical MMP on W i.e. contracting -1 -curves to get a minimal S . As it is well known S is $\mathbb{P}^2$ or a smooth ruled surface with no -1 -curves.

Let $B_S^+ = \sum b_{i,S}^+ B_{i,S}^+$ be the pushdown of B_W^+ on S where $K_W + B_W^+$ is the crepant pullback of $K_X + B^+$. Then define

$$A_S := \frac{b_{1,S}^+}{2} B_{1,S}^+ + \sum_{i \neq 1} b_{i,S}^+ B_{i,S}^+$$

If $S = \mathbb{P}^2$ then $-(K_S + A_S)$ is ample and $\text{Supp } A_S = \text{Supp } B_S^+$. By lemma 1.7.6 $(S, \text{Supp } A_S = \text{Supp } B_S^+)$ is bounded. Then lemma 1.7.7 implies the boundedness of W and so of X .

Now assume that S is a ruled surface. If there is no exceptional curve (with negative self-intersection) on S then $-(K_S + A_S)$ is nef and big if we take $B_{1,S}^+$ a non-fibre component of B_S^+ . Since S is smooth then S is bounded and so $(S, \text{Supp } A_S = \text{Supp } B_S^+)$ is bounded.

But if there is an exceptional divisor C on S then contract C as $S \rightarrow S'$. So S is a minimal resolution of S' . Since $\rho(S) = 2$ and $(S', 0)$ is δ -lc then the index of each integral divisor on S' is bounded. So S' is bounded and then $(S', \text{Supp } B_{S'}^+)$ is also bounded. This implies the boundedness of S , W and so of X . Note that $B_{S'}^+ \neq 0$ as S' is WLF.

□

Corollary 1.7.10 *Conjecture 1.1.2 holds for any finite set $\Gamma_f \subseteq [0, 1]$ of rational numbers.*

Proof It follows from corollary 1.7.9 $\square$.

1.8 Second proof of the global case

Remember that all the varieties are algebraic surfaces unless otherwise stated. We first prove the boundedness of varieties and then prove the boundedness of complements. This is somehow the opposite of what we did in the last subsection. However our proof was inspired by the theory of complements. The following proof heavily uses the properties of surfaces. That means that it is not expected to have a higher dimensional generalisation. The method also has some similarity with the proof of Alexeev and Mori [AM] in the sense that we both analyse a series of blow ups, but in different ways.

Theorem 1.8.1 *The $BAB_{\delta,2,[0,1]}$ holds.*

Proof Now we reduce to the case $B = 0$. Run the anti LMMP on the pair $(X, 0)$ i.e. if $-K_X$ is not nef then contract an extremal ray R where $K_X \cdot R > 0$. This obviously contracts a curve in B . Repeating this process gives us a model $(X', 0)$ where $-K_{X'}$ is nef and big. Otherwise X' should be with Picard number one and $K_{X'}$ nef. But this is impossible by our assumptions. We will prove the boundedness of $\{X'\}$ and so it implies the boundedness of $\{X\}$. Now we replace (X, B) with $(X', 0)$ but we denote it as $(X, 0)$. We also assume that $\delta < 1$ otherwise X will be smooth and so with bounded index.

Let $W \rightarrow X$ be a minimal resolution. The main idea is to prove that there are only a bounded number of possibilities for the coefficients in B_W where $K_W + B_W = {}^*K_X$ i.e. the index of K_X is bounded.

Strategy: Here we again have the familiar division into non-exceptional and exceptional cases.

First assume that $(X, 0)$ is non-exceptional. So there will be a $(0, n)$ -complement $K_X + B^+$ for $n < 58$. If we run the classical MMP on the pair $(W, 0)$ then we end up with S which is either $\mathbb{P}^2$ or a ruled surface. Since $-(K_S + B_S) = -{}^*(K_W + B_W)$ is nef and big then K_S can not be nef. Let $K_W + B_W^+ = {}^*(K_X + B_X^+)$

Lemma 1.8.2 *Let G be a component in the boundary B_S^+ where $K_S + B_S^+ = {}^*(K_W + B_W^+)$ then G^2 is bounded from below and above. Moreover there are only a bounded number of components in B_S^+ .*

Proof The boundedness of G^2 follows from the next lemma and the fact that X is δ -lc. The boundedness of number of components in B_S^+ is left to the reader. $\square$

The more general lemma below will also be needed later.

Lemma 1.8.3 *Let (T, B_T) be an δ -lc WLF pair where T is either $\mathbb{P}^2$ or a smooth ruled surface (with no -1 -curves) and suppose $K_T + \overline{B}$ is antinef (lc) for a boundary $\overline{B}$. Let M, B'_T be effective divisors with no common component such that $\overline{B} = B'_T + M$ then M^2 is bounded from above.*

Proof First assume that $T = \mathbb{P}^2$. In this case the lemma is obvious because if M^2 is too big then so is $\deg M$ and so it contradicts the fact that $\deg M \leq 3$.

Now assume that T is a ruled surface where F is a general fibre other than those curves in the boundary and C a section. The Mori cone of T is generated by its two edges. F generates one of the edges. If all the components of M are fibres then $M^2 = 0$ and we are done. So assume otherwise and let $M \equiv aC + bF$ then $0 < M.F = (aC + bF).F = a$ so a is positive. Let $C^2 = -e$ and consider the following two cases:

1. $e \geq 0$: We know that $K_T \equiv -2C + (2g - 2 - e)F$ where g is a non-negative number [H, V, 2.11]. So we have

$$0 \geq (K_T + M + tC).F = -2 + a + t$$

for some $t \geq 0$ where $B'_T \equiv tC + uF$ ($u \geq 0$ since $e \geq 0$). Hence $a + t \leq 2$. Calculations give $M^2 = a(2b - ae)$. Since a and e are both non-negative then M^2 big implies that b is big. But on other hand we have:

$$0 \geq (K_T + M + tC).C = (-2 + a + t)(-e) + 2g - 2 - e + b$$

This gives a contradiction if b is too big because e is also bounded. The boundedness of e follows from the fact that T is δ -lc. In order in the local isomorphic subsection we proved that exceptional divisors have bounded selfintersection numbers.

2. $e < 0$: in this case by [H, V, 2.12] we have $e + 2g \geq 0$ and so:

$$\begin{aligned} 0 &\geq (K_T + M).C = \\ &(-2 + a)(-e) + 2g - 2 - e + b = 2g + e - 2 - (ae/2) + (2b - ae)/2 \end{aligned}$$

Now since $2g + e - (ae/2) \geq 0$ then $(2b - ae)/2 \leq 2$. So M^2 is bounded because a is also bounded. $\square$

Let $P \in X$ be a singular point. If P is not in the support of B^+ then the index of K_X at P is at most 57 and so bounded. Now suppose that P is in the support of B^+ . If the singularity of P is of type E_6 , E_7 , E_8 or D_r then again the index of K_X at P is bounded. So assume that the singularity at P is of type A_r . The goal is to prove that the number of curves in $\text{exc}(W/P)$ is bounded. In order we should prove that the number of -2 -curves are bounded because the number of other curves is bounded by the proof of local isomorphic case. Note that the coefficient of any $E \in \text{exc}(W/P)$ in B_W^+ is positive and there are only a bounded number of possibilities for these coefficients. Let $\mathcal{C}$ be the longest connected subchain of -2 -curves in $\text{exc}(W/P)$. Run the classical MMP on W to get a model W' such that there is a -1 -curve F on W' s.t. it is the first -1 -curve that intersects the chain $\mathcal{C}$ (if there is no such W' and F then $\mathcal{C}$ should consist of a single curve). We have two cases:

1. F intersects, transversally and in one point, only one curve in $\mathcal{C}$ say E . First suppose that E is a middle curve i.e. there are E' and E'' in the chain which both intersect E . Now contract F so E becomes a -1 -curve. Then contract E and then E' and then all those which are on the side of E' . In this case by contracting each curve we increase E''^2 by one. And so E'' will be a divisor on S in B_S^+ with high self-intersection. By the lemma above there can be only a bounded number of curves in $\mathcal{C}$ on the side of E' . Similarly there are only a bounded number of curves on the side of E'' . So we are done in this case.

Now suppose that E is on the edge of the chain and intersects E' . Let t_E and t_F be the coefficients of E and F in B_W^+ and similarly for other curves. Let h be the intersection number of F with the curves in $B_{W'}^+$ except those in $\mathcal{C}$ and F itself. Now we have

$$0 = (K_{W'} + B_{W'}^+).F = t_E + h - 1 - t_F$$

So $h = 1 + t_F - t_E$. If $h \neq 0$ then F intersects some other curve not in the chain $\mathcal{C}$. By contracting F then E and then other curves in the chain we get a contradiction again. Now suppose $h = 0$ i.e. $t_E = 1$ and $t_F = 0$. In this case let x be the intersection of E with the curves in $B_{W'}^+$ except those in $\mathcal{C}$. So we have

$$0 = (K_{W'} + B_{W'}^+).E = -2t_E + t_{E'} + x$$

So $x = 2t_E - t_{E'} > 0$ and similarly we again get a contradiction.

2. Now assume that F intersects the chain in more than one curve or intersects a curve with intersection number more than one. Suppose the chain $\mathcal{C}$ consists of $E_1, \dots, E_s$ and F intersects $E_{j_1}, \dots, E_{j_l}$. Note that l is bounded. If $F.E_{j_k} > 1$ for all $0 \leq k \leq l$ then contract F . So $E_{j_k}^2 \geq 0$ after contraction of F and they will not be contracted later and so they appear in the boundary B_S^+ . Now replace $\mathcal{C}$ with longest connected subchain when we disregard all E_{j_k} . Now go to step one again and if it doesn't hold come back to step two and so on.

Now suppose $F.E_{j_k} = 1$ for some k . So F should intersect at least another E_{j_t} where $t = k + 1$ or $t = k - 1$. Now contract F so E_{j_k} becomes a -1 -curve and would intersect E_{j_t} . Contracting E_{j_k} and possible subsequent -1 -curves will prove that there are a bounded number of curves between E_{j_t} and E_{j_k} . Now after contracting E_{j_k} and all other curves between E_{j_t} and E_{j_k} we will have $E_{j_m}^2 \geq 0$ for each $m \neq k$. So again we take the longest connected subchain excluding all E_{j_t} . And repeat the procedure. It should stop after a bounded number of steps because the number of curves in B_S^+ is bounded. This boundedness implies that there are only a bounded number of possibilities for the coefficients in B_W where $K_W + B_W = *K_X$. By Borisov-McKernan W belongs to a bounded family and so complements would be bounded.

Here the proof of the non-exceptional case finishes and from now on we assume that $(X, 0)$ is exceptional.

Let $W \rightarrow X$ be a minimal resolution. Let $0 < \tau < \frac{1}{2}$ be a number and the minimal log discrepancy of $(X, 0)$ be $a = \text{mld}(X, 0)$. If $a \geq \frac{1}{2} + \tau$ then we know that X belongs to a bounded family according to step 1 in the proof of theorem 1.7.1 above. So we assume $a < \frac{1}{2} + \tau$ and then blow up all exceptional/ X curves E with log discrepancy $a_E = a(E, X, 0) \leq \frac{1}{2} + \tau$ to get $Y \rightarrow X$ and put $K_Y + B_Y = *K_X$. Fix E_1 , one of these exceptional divisors. Let $t \geq 0$ be a number such that there is an extremal ray R such that $(K_Y + B_Y + tE_1).R = 0$ and $E_1.R > 0$ (and s.t. $K_Y + B_Y + tE_1$ is Klt and antinef). Such R exists otherwise there is a $t > 0$ such that $K_Y + B_Y + tE_1$ is lc (and not Klt) and antinef. This is a contradiction by [Sh2, 2.3.1]. Now contract $R : Y \rightarrow Y_1$ if it is of birational type.

Again by increasing t there will be an extremal ray R_1 on Y_1 such that $(K_{Y_1} + B_{Y_1} + tE_1).R_1 = 0$ and $E_1.R_1 > 0$ (preseving the nefness of $-(K_{Y_1} + B_{Y_1} + tE_1)$). If it is of birational type then contract it and so on. After finitely many steps we get a model $(V_1, B_{V_1} + t_1E_1)$ and a number $t_1 > 0$

with the following possible outcomes:

(1.8.3.1)

- ◇ $(V_1, B_{V_1} + t_1 E_1)$ is Klt, $\rho(V_1) = 1$ and $K_{V_1} + B_{V_1} + t_1 E_1 \equiv 0$.
- ◇ $(V_1, B_{V_1} + t_1 E_1)$ is Klt and $\rho(V_1) = 2$ and there is a non-birational extremal ray R on V_1 such that $(K_{V_1} + B_{V_1} + t_1 E_1).R = 0$. Moreover $K_{V_1} + B_{V_1} + t_1 E_1$ is antinef.

Note that for each element $E \in \text{exc}(Y/X)$, either E is a divisor on V_1 or it is contracted to a point in the support of E_1 .

Lemma 1.8.4 *For any $h > 0$ there is an $\eta > 0$ such that if (T, B) is a δ -lc pair (δ is already fixed) with a component C of B passing through $P \in T$, with a coefficient $t \geq h$, then either K_T is $\delta + \eta$ -lc at P or $1 - a_E > \eta$ for each exceptional divisor E/P on a minimal resolution of T ($a_E = \log$ discrepancy of (T, B) at E).*

Proof If P is smooth or has E_6, E_7, E_8 or D_r type of singularity then the lemma is clear since the index of K_T at P is bounded in all these cases (see the local isomorphism subsection). In order in all these cases there will be an $\eta > 0$ such that K_T is $\delta + \eta$ -lc at P .

Now suppose that the singularity at P is of type A_r . Take a minimal resolution $W_T \rightarrow T$ with $\text{exc}(W_T/P) = \{E_1, \dots, E_r\}$ (notation as in the local isomorphism subsection) and suppose that j is the maximal number such that $\text{mld}(P, T, 0) = a'_j$ (here we show the log discrepancy of $(T, 0)$ at E_* as a'_*) for an exceptional divisor E_j/P . Actually we may assume that $r - j$ is bounded. By the local isomorphism subsection, the distance of E_j from one of the edges of $\text{exc}(W_T/P)$ is bounded. We denote the birational transform of C on W_T again by C . Suppose C intersects E_k in $\text{exc}(W_T/P)$. If $k \neq 1$ or r then we have $(-E_k^2)a_k - a_{k-1} - a_{k+1} + x = 0$ where a_* shows the log discrepancy of the pair (T, B) at E_* and $x \geq h$ a number. So either $a_{k-1} - a_k \geq \frac{h}{2}$ or $a_{k+1} - a_k \geq \frac{h}{2}$. In either case the distance of E_k is bounded from one of the edges of $\text{exc}(W_T/P)$. If this edge is the same edge as for E_j then again the lemma is clear since the coefficients of E_k and E_j in $*C$ (now C is on T and $*C$ on W_T) are bounded from below (in other words they are not too small). Now assume the otherwise i.e. E_k and E_j are close to different edges. In this case we claim that the coefficients of the members of $\text{exc}(W_T/P)$ in

$\overline{B}_{W_T}$, where $K_W + \overline{B}_{W_T} = *(K_T + tC)$, are bounded from below. Suppose that the smallest coefficient occurs at E_m . Simple calculation shows that we can assume that E_m is one of the edges of $\text{exc}(W_T/P)$. Hence E_m is with a bounded distance from E_j or from E_k .

Suppose that E_m is with a bounded distance from E_j . If $a'_j \geq \frac{1+\delta}{2}$ then K_T is $\frac{1+\delta}{2}$ -lc at P . So we can assume that $a'_j < \frac{1+\delta}{2}$. We prove that all the numbers $1 - a'_j, \dots, 1 - a'_r$ are bounded from below. In order, if $1 < j < r$ then $(-E_j^2)a'_j - a'_{j-1} - a'_{j+1} = 0$ (note that $-E_j^2 > 2$ in this case). Now if $a'_{j-1} - a'_j \geq \frac{\delta}{2}$ then the chain will be bounded and so the index of K_T at P . But if $a'_{j+1} - a'_j \geq \frac{\delta}{2}$ then $a'_r - a'_{r-1} \geq \frac{\delta}{2}$ and so $(-E_r^2 - 1)a'_r = 1 - (a'_r - a'_{r-1}) \leq 1 - \frac{\delta}{2}$. Hence if $m = r$ then we are done. But if $m = 1$ then again the whole chain is bounded and so the index of K_T at P . Now if $j = r$ then again the chain is bounded if $m = 1$ and $a'_m = a'_j = a'_r < \frac{1+\delta}{2}$ if $m = r$.

In the second case i.e. if E_m is with a bounded distance from E_k then the coefficient of E_m in $*C$ on W is bounded from below. $\square$

Lemma 1.8.5 *For any $h > 0$ there is a $\gamma > 0$ such that if (T, B) is a δ -lc pair (δ is already fixed), WLF with a component C of B passing through $P \in T$ and $t \geq h$ where t is the coefficient of C in B , then K_T is $\delta + \gamma$ -lc.*

Proof As discussed in lemma 1.8.4 we may assume that the singularity at P is of type A_r and $1 - a_k > \eta$ for some fixed number $\eta > 0$ where a_k is the log discrepancy of the pair (T, B) at any exceptional divisor E_k/P on W_T where $W_T \rightarrow T$ is a minimal resolution (we put $\text{exc}(W_T/P) = \{E_1, \dots, E_r\}$). Let $\mathcal{C}$ be the longest connected sub-chain of -2 -curves in $\text{exc}(W_T/P)$ and W_1 a model where $\mathcal{C}$ is intersected by a -1 -curve F for the first time i.e. we blow down -1 -curves on W_T till we get a model W_1 and a morphism $W_T \rightarrow W_1$ such that W_1 is the first model where there is a -1 -curve F intersecting $\mathcal{C}$ (on W_1). Let $K_{W_T} + B^+ \equiv 0$ be a (lc) $\mathbb{Q}$ -complement of $K_{W_T} + B_{W_T}$. Assume that F intersects E_j in $\mathcal{C}$ and let t_{E_j} and t_F be the coefficients of E_j and F in B^+ on W_T (similar notation for the coefficients of other exceptional divisors). Then an argument as in the proof of the non-exceptional case gives a contradiction:

1. Suppose F intersects, transversally and in one point, only one curve in $\mathcal{C}$ (which is E_j). First suppose that E_j is a middle curve i.e. there are E_{j-1} and E_{j+1} in $\mathcal{C}$ which both intersect E_j . Now contract F so E_j becomes a -1 -curve. Then contract E_j and then E_{j-1} and then all those which are

on the of E_{j-1} . By contracting each curve we increase E_{j+1}^2 by one. If we continue contracting -1 -curves we get S ($S = \mathbb{P}^2$ or a ruled surface with no -1 -curve) where E_{j+1} is a component of B_S . By lemma 1.8.3 there can be only a bounded number of curves in $\mathcal{C}$ on the side of E_{j-1} . Similarly there are only a bounded number of curves in $\mathcal{C}$ on the side of E_{j+1} . So we are done in this case.

Now suppose that E_j is on the edge of the chain $\mathcal{C}$ and it intersects E_{j-1} . Let $B^+_{W_1} = B^+ + M$ (M and B^+ with no common component) where each component of B^+ is either F or an element of $\mathcal{C}$. Now we have

$$0 = (K_{W_1} + B^+_{W_1}).F = t_{E_j} - 1 - t_F + (M.F)$$

So $M.F = 1 + t_F - t_{E_j}$. Similarly let $B^+_{W_1} = \ddot{B}^+ + N$ (N and $\ddot{B}^+$ with no common component) where each component of $\ddot{B}^+$ is either F or an element of $\mathcal{C}$. Then we have

$$0 = (K_{W_1} + B^+_{W_1}).E_j = -2t_{E_j} + t_{E_{j-1}} + t_F + (N.E_j)$$

and so $t_{E_j} = t_{E_{j-1}} - t_{E_j} + t_F + (N.E_j) > \eta$. Hence $t_{E_{j-1}} - t_{E_j} > \frac{\eta}{3}$ or $t_F > \frac{\eta}{3}$ or $(N.E_j) > \frac{\eta}{3}$.

If $t_F > \frac{\eta}{3}$ then by contracting F we increase M^2 at least by $(M.F)^2 \geq t_F^2 > (\frac{\eta}{3})^2$. We have the same increase when we contract E_j and then E_{j-1} and so on. So lemma 1.8.3 shows the boundedness of $\mathcal{C}$.

If $(N.E_j) > \frac{\eta}{3}$ then proceed similar to the last paragraph.

If $t_{E_{j-1}} - t_{E_j} > \frac{\eta}{3}$ then $t_{E_{j-1}} > t_{E_j} + \frac{\eta}{3}$. This implies that $t_{E_j} \leq 1 - \frac{\eta}{3}$ then $M.F \geq \frac{\eta}{3}$ and so we continue as above.

2. Now assume that F intersects $\mathcal{C}$ in more than one curve or intersects a curve in $\mathcal{C}$ with intersection number more than one. Suppose the chain $\mathcal{C}$ consists of $E_s, \dots, E_u$ and F intersects $E_{j_1}, \dots, E_{j_l}$. Note that l is bounded.

If $F.E_{j_k} > 1$ for all $1 \leq k \leq l$ then contract F . So $E_{j_k}^2 \geq 0$ after contraction of F and hence E_{j_k} can not be contracted and so it appears in the boundary on a “minimal” model S (i.e. S is the projective plane or a smooth ruled surface with no -1 -curve). Replace $\mathcal{C}$ with its longest connected subchain when we disregard all E_{j_k} . From here we can go back to step one and repeat the argument.

Now suppose $F.E_{j_k} = 1$ for some k . So F should intersect at least another E_{j_q} where $q = k+1$ or $q = k-1$. Now contract F so E_{j_k} becomes a -1 -curve and would intersect E_{j_q} . Contracting E_{j_k} and possible subsequent -1 -curves will prove that there are only a bounded number of curves between E_{j_q} and

E_{j_k} in $\mathcal{C}$. Now after contracting E_{j_k} and all other curves between E_{j_q} and E_{j_k} we will have $E_{j_m}^2 \geq 0$ for each $m \neq k$. So again we take the longest connected subchain excluding $E_{j_1}, \dots, E_{j_l}$ and go back to step one.

This process should stop after a bounded number of steps because the number of curves in B_S^+ with coefficient $> \eta$ is bounded (S is again a “minimal” model). To prove this later boundedness note that $(K_S + B_S^+).F = 0$, where we assume that S is a ruled surface and F a fibre, implies that there are only a bounded number of non-fibre components in B_S^+ with coefficient $> \eta$. Let L be a section and t_L be its coefficient in B_S^+ and F_i fibre components of B_S^+ with $t_{F_i} > \eta$. So

$$\begin{aligned} 0 &\geq (K_S + t_L L + \sum_i t_{F_i} F_i).L = (-2L + (2g - 2 - e)F + t_L L + \sum_i t_{F_i} F_i).L \\ &= -t_L e + e + 2g - 2 + \sum_i t_{F_i} \end{aligned}$$

which proves that there are a bounded number of F_i ($L^2 = -e$ and $e + 2g \geq 0$ if $e < 0$). So the chain $\mathcal{C}$ should have a bounded length. This implies that if we throw C away in the boundary B then the mld at P will increase at least by $\gamma > 0$ a fixed number (which doesn't depend on P nor T). This proves the lemma. $\square$

Lemma 1.8.5 settles the first case in 1.8.3.1 by deleting the boundary B_{V_1} . Now assume the second case in the division above in 1.8.3.1. Let F be a general fibre of the contraction defined by the extremal ray R . If the other extremal ray of V_1 defines a birational map $V_1 \rightarrow Z$ (otherwise delete the boundary and use 1.8.5) then let H be the exceptional divisor of this contraction.

If K_{V_1} is antinef then again use 1.8.5. If K_{V_1} is not antinef and if $E_1 \neq H$ then apply lemma 1.8.5 to (Z, B_Z) . Boundedness of Z implies the boundedness of V_1 and so we can apply lemma 1.7.6. But if K_{V_1} is not antinef and $E_1 = H$ then perform a hat of the third type as defined in the proof of theorem 1.7.1 where $(U, G_U) := (V_1, B_{V_1} + t_1 E_1)$ and $V_2 := U'$. We can use lemma 1.8.5 on V_2 or after contracting a curve on V_2 , in order, to get the boundedness of V_2 . Boundedness of V_2 implies the boundedness of V_1 . $\square$

Corollary 1.8.6 *Conjecture $WC_{\delta, 2, \Gamma_f}$ holds in the global case where Γ_f is a finite subset of rational numbers in $[0, 1)$.*

Proof Obvious by theorem 1.8.1.

1.9 An example

Example 1.9.1 Let m be a positive natural number. For any $1 > \eta > 0$ and any $\tau > 0$ there is a model $(X, 0)$ (may not be global) satisfying the followings:

1. X is $\frac{1}{m}$ -lc.
2. Suppose $Y \rightarrow X$ is a partial resolution such that $K_Y + B_Y = {}^*K_X$ is $\frac{1}{m} + \eta$ -lc and $b_i > \frac{m-1}{m} - \eta$. Put $D = \sum \frac{m-1}{m} B_i$.
3. $K_Y + D$ is not $\frac{1}{m} + \tau$ -lc.

Proof Let $P \in X$ and X smooth outside P . Suppose the minimal resolution of P has the following diagram:

$$O^{-3} \text{ --- } O^{-2} \text{ --- } \dots \text{ --- } O^{-2} \text{ --- } O^{-2} \text{ --- } O^{-4}$$

where the numbers show the self-intersections.

This diagram has the following corresponding system on a minimal resolution where a_i stand for the log discrepancies:

$$\begin{aligned} 3a_1 - a_2 - 1 &= 0 \\ 2a_2 - a_1 - a_3 &= 0 \\ &\vdots \\ 2a_{r-1} - a_{r-2} - a_r &= 0 \\ 4a_r - a_{r-1} - 1 &= 0 \end{aligned}$$

Now put $a_{r-1} - a_r = t$ so $a_{r-2} - a_{r-1} = t, \dots, a_1 - a_2 = t$. So we have: $a_r = \frac{1+t}{3}$ and $a_1 = \frac{1-t}{2}$. The longer the chain is the smaller the t is and the discrepancies vary from $-\frac{1+t}{2}$ to $\frac{t-2}{3}$. Other a_i can be calculated as $a_i = a_1 - (i-1)t = \frac{1-t}{2} - (i-1)t = \frac{1-(2i-1)t}{2}$.

Suppose j is such that $a_j < \frac{1}{m} + \eta$ but $a_{j-1} \geq \frac{1}{m} + \eta$. So the exceptional divisors corresponding to $a_r, a_{r-1}, \dots, a_j$ will appear on Y but others not. Now we try to compute the log discrepancies of the pair (Y, D) . In order

the minimal resolution for $P \in X$ is also the minimal resolution for Y . But here just $E_1, \dots, E_{j-1}$ are exceptional/ Y . The system for the new log discrepancies (for (Y, D)) is as follows:

$$\begin{aligned} 3a'_1 - a'_2 - 1 &= 0 \\ 2a'_2 - a'_1 - a'_3 &= 0 \\ &\vdots \\ 2a'_{j-2} - a'_{j-3} - a'_{j-1} &= 0 \\ 2a'_{j-1} - a'_{j-2} - \frac{1}{m} &= 0 \end{aligned}$$

Again put $a'_{j-2} - a'_{j-1} = s$ so similarly we have $a'_{j-1} = \frac{1}{m} + s$ and $a'_1 = \frac{1-s}{2}$. If j is big (i.e. if t is small enough) then s would be small and so $a'_{j-1} = \frac{1}{m} + s < \frac{1}{m} + \tau$. Hence (Y, D) is not $\frac{1}{m} + \tau$ -lc. $\square$

1.10 Local cases revisited

Using the methods in the proof of the global case, we give a new proof of the local cases. Here again by $/Z$ we mean $/P \in Z$ for a fixed P . The following is the main theorem in this subsection.

Theorem 1.10.1 *Conjecture $WC_{\delta, 2, \Phi_{sm}}$ holds in the local case i.e. when we have $\dim Z \geq 1$.*

Proof Our proof is similar to the non-exceptional global case. Here the pair $(X/Z, B)$ is a relative WLF surface log pair (i.e. $-(K_X + B)$ is nef and big/ Z), where (X, B) is δ -lc and $B \in \Phi_{sm}$. Fix $P \in Z$. Then there exists a regular $(0, n)$ -complement/ $P \in Z$, $K + B^+$ for some $n \in \{1, 2, 3, 4, 6\}$ by [Sh2].

1. Remember the first step in the proof of theorem 1.7.1.
2. Remember definition 1.7.2 and lemma 1.7.3. Let m be the smallest number such that $\frac{1}{m} \leq \delta$. Let $h = \min\{\frac{k-1}{k} - \frac{u}{r!} > 0\}_{1 \leq k \leq m}$ where u, k are natural numbers and $r = \max\{m, 6\}$. Now choose a τ for m as in lemma 1.7.3 such that $\tau < h$.

Blow up one exceptional divisor E/P via $f : Y \longrightarrow X$ such that the log discrepancy satisfies $\frac{1}{k} \leq a(E, X, B) \leq \frac{1}{k} + \tau$ for some k (if such E doesn't exist then go to step 1). The crepant log divisor $K_Y + B_Y$ is $\frac{1}{m}$ -lc and so by the choice of τ , $K_Y + D_\tau$ is also $\frac{1}{m}$ -lc (D_τ is constructed for B_Y). Let $K_Y + B_Y^+$ be the crepant blow up of $K_X + B^+$. Then again by the way we chose τ we have $D_\tau \leq B_Y^+$. Now run the anti-LMMP/ $P \in Z$ over $K_Y + D_\tau$ i.e. contract any birational type extremal ray $R/P \in Z$ such that $(K_Y + D_\tau).R > 0$. At the end we get a model X_1 with one of the following properties:


- ◇ $(K_{X_1} + D_\tau) \equiv 0/P \in Z$ and $K_{X_1} + D_\tau$ is $\frac{1}{m}$ -lc.
- ◇ $-(K_{X_1} + D_\tau)$ is nef and big/ $P \in Z$ and $K_{X_1} + D_\tau$ is $\frac{1}{m}$ -lc.

where $K_{X_1} + D_\tau$ is the birational transform of $K_Y + D_\tau$ and let $g : Y \longrightarrow X_1$ be the corresponding morphism.

The nefness of $-(K_{X_1} + D_\tau)$ comes from the fact that $D_\tau \leq B_1^+$. And $K_{X_1} + D_\tau$ is $\frac{1}{m}$ -lc by applying lemma 1.7.3.

3. Whichever case occurs above, to construct a complement, it is enough to bound the index of $K_{X_1} + D_\tau/P$.
4. Let C be a curve contracted by $g : Y \longrightarrow X_1$. If C is not a component of B_Y then the log discrepancy of C with respect to $K_{X_1} + B_{X_1}$ is at least 1 where $K_{X_1} + B_{X_1}$ is the birational transform of $K_Y + B_Y$. Moreover $g(C) \in \text{Supp } B_{X_1} \neq \emptyset$. So the log discrepancy of C with respect to K_{X_1} is more than 1. This means that C is not a divisor on a minimal resolution $W_1 \longrightarrow X_1$. Let $W \longrightarrow X$ be a minimal resolution. Then there is a morphism $W \longrightarrow W_1$. Hence $\text{exc}(W_1/X_1) \subseteq \text{exc}(W/X) \cup \text{Supp}(B = B_X)$. Now if $C \in \text{exc}(W/X) \cup \text{Supp } B$ is contracted by g then $a(C, X_1, D_\tau) < a(C, X, B)$.
5. Let $(X_1, B_1) := (X_1, D_\tau)$ and repeat the process. In other words again we blow up one exceptional divisor E via $f_1 : Y_1 \longrightarrow X_1$ such that the log discrepancy satisfies $\frac{1}{k} \leq a(E, X_1, B_1) \leq \frac{1}{k} + \tau$ for some natural number $k > 1$. The crepant log divisor $K_{Y_1} + B_{1,Y_1}$ is $\frac{1}{m}$ -lc and so by lemma 1.7.3 $K_{Y_1} + D_{1,\tau}$ is $\frac{1}{m}$ -lc. Note that the point which is blown up on X_1 can not be smooth since $\tau < h$ as defined above. So according to the last step the blown up divisor E is a member of $\text{exc}(W/X) \cup \text{Supp } B$.

Now we again run the anti-LMMP on $K_{Y_1} + D_{1,\tau}$ and proceed as in step 2.



6. Steps 4,5 show that each time we blow up a member of $\text{exc}(W/X) \cup \text{Supp } B$ say E . And if we blow that divisor down in some step then the log discrepancy $a(E, X_j, B_j)$ will decrease. That divisor will not be blown up again unless the log discrepancy drops at least by $\frac{1}{2(m-1)} - \frac{1}{2m}$ (this is not a sharp bound). So after finitely many steps we get a model X_i with a standard boundary B_i such that there is no E/P where $\frac{1}{k} \leq a(E, X_i, B_i) \leq \frac{1}{k} + \tau$ for any $1 < k \leq m$. Hence the index of $-(K_{X_i} + B_i)/P$ is bounded and so we can construct an appropriate complement for $(X_i, B_i)/Z$. This implies the existence of the desired complement for $(X, B)/Z$.

□

2 ϵ -log canonical complements in higher dimensions

In this section we consider the (ϵ, n) -lc complements in higher dimensions i.e. in dimensions more than two. This is a joint work in progress with V.V. Shokurov. In subsection 2.1 we try to work out the proof of theorem 1.7.1 in dim 3 and we point out the problems we have to solve in order to finish the proof of conjecture 1.1.2 in dim 3 (this is the plan of the author). In subsection 2.2 we outline Shokurov's plan on the same problem. These plans have already won an EPSRC three years postdoctoral fellowship by the author.

Let $X \rightarrow Z$ be an extremal K_X -negative contraction where X is a 2-dim Pseudo-WLF and Z is a curve. We know that $Z \simeq \mathbb{P}^1$ since Z should be rationally connected as X is. Moreover $\rho(X) = 2$. Similar Mori fibre spaces in higher dimensions are not that simple. This makes the boundedness problem of (ϵ, n) -lc complements more difficult in higher dimensions. We also don't know yet whether the index of $K_X + B$ will be bounded if we fix the mld at a point.

In section 1 we first proved the boundedness of ϵ -lc complements and then the BAB. But in higher dimensions we expect to prove both problems together at once. In other words in some cases where it is difficult to prove the boundedness of varieties, it seems easier to prove the boundedness of complements; specially when we deal with a fibre space. Conversely when it is difficult to prove the boundedness of ϵ -lc complements, it is better to prove the boundedness of pairs; this is usually the case when the pairs are exceptional.

Lemma 2.0.2 *Let $X \dashrightarrow X'$ be a flip/ Z and assume that (X, B) is (ϵ, n) -complementary/ Z then (X', B') is (ϵ, n) -complementary/ Z where B' is the birational transform of B .*

Proof Obvious from the definition of (ϵ, n) -complements.

Note that in the previous lemma it doesn't matter that the flipping is with respect to which log divisor.

Lemma 2.0.3 *Let (Y, B) be a pair and $Y \dashrightarrow Y'/Z$ be a composition of divisorial contractions and flips/ Z such that in each step we contract an*

extremal ray R where $(K+B).R \geq 0$. Suppose $B' = \sum b'_i B'_i$ is the birational transform of B , the pair (Y', B') is (ϵ, n) -complementary/ Z and $(n+1)b'_i \geq nb'_i$ for each coefficient b'_i then (Y, B) is also (ϵ, n) -complementary/ Z .

Proof Clear by lemmas 1.2.10 and 2.0.2.

Lemma 2.0.4 *The Klt Pseudo-WLF property is preserved under extremal flips and divisorial contractions with respect to any log divisor.*

Proof Let X be a Klt Pseudo-WLF and B a boundary such that (X, B) is a Klt WLF. Now let $X \dashrightarrow X'$ be an extremal flip corresponding to an extremal ray R . Since (X, B) is a Klt WLF then there is a rational boundary D such that $K_X + D$ is antiample and Klt. Now let H' be an ample divisor on X' and H its transform on X . There is a rational $t > 0$ such that $K_X + D + tH$ is antiample and Klt. Now take a Klt $\mathbb{Q}$ -complement $K_X + D + tH + A \equiv 0$. So we have $K_{X'} + D' + tH' + A' \equiv 0$ on X' . From the assumptions $K_{X'} + D' + A'$ is antiample and Klt. So X' is also a Klt Pseudo-WLF.

If $X \rightarrow X'$ is a divisorial extremal contraction then proceed as in the flip case by taking an ample divisor H' on X' .

□

Definition 2.0.5 Let (V, B_V) and (U, B_U) be lc pairs. U is called a semi-partial resolution of V if there is a partial resolution (W, B_W) of (V, B_V) such that W and U are isomorphic in codim 1.

Definition 2.0.6 (D -LMMP) Let D be an $\mathbb{R}$ -Cartier divisor on a normal variety X . We say D -LMMP holds if the followings hold:

- ◇ Any D -negative extremal ray R on X can be contracted. And the same holds in the subsequent steps for the birational transform of D .
- ◇ If a D -contraction as in the first step is a flipping then the corresponding D -flip exists.
- ◇ Any sequence of D -flips terminates.

If $D := K + B$ for a lc $\mathbb{R}$ -Cartier divisor $K + B$ then we know that D -LMMP holds in dim 3 by [Sh5].

Remark 2.0.7 Let D be an $\mathbb{R}$ -Cartier divisor on a variety X of dim d and assume that LMMP holds in dim d . Moreover assume that $\beta D \equiv K + B$ for a lc $\mathbb{R}$ -Cartier log divisor $K + B$ and $\beta > 0$ then the D -LMMP holds. Since in this case D -LMMP and $K + B$ -LMMP are equivalent.

Example 2.0.8 Let (X, B) be a d -dim Klt WLF and suppose LMMP holds in dim d then $-K$ -LMMP holds. In order since (X, B) is a Klt WLF then there is a Klt $\mathbb{Q}$ -complement $K + B^+ \equiv 0$. There is a $t > 0$ such that $K + B^+ + tB^+ \equiv tB^+$ is Klt. Since $-K \equiv B^+$ then $-K$ -LMMP is equivalent to B^+ -LMMP and again equivalent to tB^+ -LMMP. Since $K + B^+ + tB^+$ -LMMP holds so does $-K$ -LMMP.

2.1 ϵ -lc complements in dimension 3

In this subsection we propose a plan toward the resolution of conjecture 1.1.2 in dim 3.

We repeat the proof of 1.7.1, in dim 3, step by step:

1. Under the assumptions of conjecture 1.1.2 for $d = 3$ and $\Gamma = \{0\}$, first assume that $(X, 0)$ is non-exceptional.
2. We don't have much information about the accumulation points of mlds in dim 3. Actually we still have not proved ACC in dim 3. As pointed out in the introduction in section 1, only one case of ACC in dim 3 is remained to be proved. Remember that Shokurov's program tries to use complements in dim $d - 1$ to prove the ACC in dim d . So it is reasonable to assume ACC in dim $d - 1$.

Lets show the set of accumulation points of mlds of d -dim lc pairs (T, B) , where $B \in \Gamma$, with $Accum_{d, \Gamma}$.

3. We need the inductive version of complements; since $(X, 0)$ is not exceptional then it is expected that there is an inductive $(0, n)$ -complement $K_X + B^+$ where $n \in \mathcal{N}_2$.
4. Remeber definition 1.7.2. We can similarly define $D_{\tau, A}$ for a boundary B , with respect to a real number $\tau \geq 0$ and a set $A \subseteq [0, 1]$:

$$D_{\tau, A} := \sum_{b_i \notin [a-\tau, a]} b_i B_i + \sum_{b_i \in [a-\tau, a]} a B_i$$

where in the first term $b_i \notin [a - \tau, a]$ for any $a \in A$ but in the second term $a \in A$ is the biggest number satisfying $b_i \in [a - \tau, a]$.

Definition 2.1.1 Let $A \subseteq [0, 1]$ and (T, B) a log pair. We say that (T, B) is A -lc if (T, B) is x -lc where $x := 1 - \sup\{A\}$.

Assuming the ACC in dim 3 a statement similar to lemma 1.7.3 may hold: For any $\gamma > 0$ and finite set $A \subseteq [0, 1]$ containing $1 - \gamma$ there is a real number $\tau > 0$ such that if (T, B_T) is a 3-fold log pair, $P \in T$, $K_T + B_T$ is γ -lc in codim 2 at P and $D_{\tau, A} \in A$ then $K_T + D_{\tau, A}$ is also γ -lc in codim 2 at P .

Moreover we expect to choose a $\tau > 0$ such that the followings hold as well:

- If $B_T \in A$ and E the exceptional divisor of a smooth blow up of T then $a(E, T, B_T) \notin [1 - a, 1 - a + \tau]$ for any $a \in A$.
 - If $B_T \in A$ and the pair (T, B_T) non-exceptional then we can refine $\mathcal{N}_2$ such that there is a $(0, n)$ -complement $K_T + B_T^+$ for some $n \in \mathcal{N}_2$ where $B_T \leq B_T^+$.
5. Let $A_1 := \{a_1\}$ where $1 - a_1 = \max \text{Accum}_{3, \{0\}} \cap [0, \delta]$. Now blow up all exceptional divisor E such that $a(E, T, B_T) \in [1 - a, 1 - a + \tau]$ for some $a \in A_1$ to get $f : Y \rightarrow X$. Construct D_{τ, A_1} for B_Y where $K_Y + B_Y$ is the crepant pull back. So (Y, D_{τ, A_1}) is A_1 -lc. Run the D -LMMP where $D := -(K_Y + D_{\tau, A_1})$. At the end we get $Y \dashrightarrow X_1$ and $X_1 \dashrightarrow S_1$ such that $-(K_{X_1} + D_{\tau, A_1})$ is nef and $\equiv 0/S_1$ and $-(K_{S_1} + D_{\tau, A_1}) \cdot R > 0$ for any birational type extremal ray R .
 6. There are the following possibilities for the model S_1 :
 - ◊ $\rho(S_1) = 1$, $-(K_{S_1} + D_{\tau, A_1}) = -(K_{S_1} + B^+) \equiv 0$ and $K_{S_1} + D_{\tau, A_1}$ is A_1 -lc.
 - ◊ There is a fibration type extremal ray R such that $-(K_{S_1} + D_{\tau, A_1}) \cdot R = 0$ and $K_{S_1} + D_{\tau, A_1}$ is A_1 -lc.
 - ◊ $-(K_{S_1} + D_{\tau, A_1})$ is nef and big and $K_{S_1} + D_{\tau, A_1}$ is A_1 -lc.
 7. In the first case in the division above we are done. In the second and third case then replace $(X, 0)$ by $(X_1, B_1) := (X_1, D_{\tau, A_1})$ and go back

to step one and repeat. By repeating and repeating the process, each time we get new coefficients. In other words we need to replace A_i with A_{i+1} such that $A_i \subseteq A_{i+1}$. We need to prove that $\cup_{i \rightarrow \infty} A_i$ is finite.

8. At the end we get a model (X_r, B_r) which is terminal in codim 2. Then we hope to prove the boundedness of the index of $K_{X_r} + B_r$ possibly after some more blow ups and blow downs. This will settle the problem if $-(K_{X_r} + B_r)$ is nef and big. Otherwise we may have a fibration and $K_{X_r} + B_r^+ = K_{X_r} + B_r + N$ where N is vertical. Then we may replace N by N' and construct a desirable complement $K_{X_r} + B_r + N'$. At the end we need to prove that the boundedness of the complement implies the boundedness of the pairs.
9. Now let $(X, 0)$ be exceptional. Since $BAB_{1,3,\{0\}}$ holds by [KMMT] then assuming ACC in dim 3, there is a $\tau > 0$ such that $BAB_{1-\tau,3,\{0\}}$ also holds. Blow up an exceptional/ X divisor E_1 with log discrepancy $a_{E_1} = a(E_1, X, 0) \leq 1 - \tau$ to get $Y \rightarrow X$ and put $K_Y + B_Y = {}^*K_X$. Let $t \geq 0$ be a number such that there is an extremal ray R such that $(K_Y + B_Y + tE_1).R = 0$ and $E_1.R > 0$ (and s.t. $K_Y + B_Y + tE_1$ Klt and antinef). Such R exists otherwise there is a $t > 0$ such that $K_Y + B_Y + tE_1$ is lc (and not Klt) and antiample. This is a contradiction with the fact that $(X, 0)$ is exceptional. Now contract $R : Y \rightarrow Y_1$ if it is of birational type (and perform the flip if it is a flipping).

Again by increasing t there will be an extremal ray R_1 on Y_1 such that $(K_{Y_1} + B_{Y_1} + tE_1).R_1 = 0$ and $E_1.R_1 > 0$ (preserving the nefness of $-(K_{Y_1} + B_{Y_1} + tE_1)$). If it is of birational type then contract it and so on. After finitely many steps we get a model $(V_1, B_{V_1} + t_1E_1)$ and a number $t_1 > 0$ with the following possible outcomes:

- ◇ $(V_1, B_{V_1} + t_1E_1)$ is Klt, $\rho(V_1) = 1$ and $K_{V_1} + B_{V_1} + t_1E_1 \equiv 0$.
- ◇ $(V_1, B_{V_1} + t_1E_1)$ is Klt and there is a fibre type extremal ray R on V_1 such that $(K_{V_1} + B_{V_1} + t_1E_1).R = 0$ and $K_{V_1} + B_{V_1} + t_1E_1$ is antinef.

If the second case occurs then we don't know $\rho(V_1)$ unlike the surface case where $\rho(V_1) = 2$.

10. In the proof of theorem 1.7.1 we introduced three types of hat. Here also we can similarly define hats but it is not clear yet how to proceed.

2.2 ϵ -lc complements in dimension 3: Shokurov's approach

Here we explain Shokurov's approach to the problem discussed in 4.1.

1. We know that the $BAB_{1,3,\{0\}}$ holds by [KMMT]. Let a be the smallest positive real number with the following property: $BAB_{a',3,\{0\}}$ holds for any $a' > a$. The idea is to prove that $BAB_{a,3,\{0\}}$ holds and so assuming the ACC in dim 3 we can prove that $a = 0$. Now assume that $BAB_{\epsilon',3,\{0\}}$ holds for any $\epsilon' > \epsilon$ where $1 > \epsilon > 0$.
2. Prove $SC_{\epsilon,3}$ in the local case. Moreover prove that the local ϵ -lc complement indexes can be chosen such that there is a $\tau > 0$ s.t. if $1 - \epsilon - \tau \leq b \leq 1 - \epsilon$ then $\lfloor (n+1)b \rfloor \geq n(1 - \epsilon)$ for any local ϵ -lc complement index n .
3. Blow up all exceptional divisor E such that $\epsilon \leq a(E, X, 0) \leq \epsilon + \tau$ to get $f : Y \rightarrow X$. Then $D_{\tau,\{1-\epsilon\}} := \sum_i (1 - \epsilon)B_i$ where $B_Y = \sum_i b_i B_i$ is the crepant pull back boundary. Then run the D -LMMP for $D := -(K_Y + D_{\tau,\{1-\epsilon\}})$. At the end we get $g : Y \dashrightarrow X_1$ and $X_1 \dashrightarrow S_1$ such that $-(K_{X_1} + D_{\tau,\{1-\epsilon\}})$ is nef and $\equiv 0/S_1$ and $-(K_{S_1} + D_{\tau,\{1-\epsilon\}}).R > 0$ for any birational type extremal ray R .
4. There are the following possibilities for the model S_1 :
 - ◇ $\rho(S_1) = 1$, $K_{S_1} + D_{\tau,\{1-\epsilon\}}$ is ample and $K_{S_1} + D_{\tau,\{1-\epsilon\}}$ is ϵ -lc.
 - ◇ $-(K_{S_1} + D_{\tau,\{1-\epsilon\}}).R = 0$ for a fibre type extremal ray R and the log divisor $K_{S_1} + D_{\tau,\{1-\epsilon\}}$ is ϵ -lc.
 - ◇ $-(K_{S_1} + D_{\tau,\{1-\epsilon\}})$ is nef and big and $K_{S_1} + D_{\tau,\{1-\epsilon\}}$ is ϵ -lc.
5. If the first case happens in the division above then delete the boundary, so $(S_1, 0)$ is $\epsilon + \tau$ -lc and so the pair is bounded by the assumptions.
6. **Definition 2.2.1** Let $f : T \rightarrow Z$ be a contraction and $K_T + B \sim_{\mathbb{R}} 0/Z$. Put $D_Z := \sum_i d_i D_i$ where d_i is defined as follows:

$$1 - d_i = \sup\{c | K_T + B + cf^*D_i \text{ is lc over the generic point of } D_i\}$$

7. If the second case occurs in the division above then we need the following general conjecture, due to Shokurov [PSh1] and Kawamata [K3], which is useful in many situations:

Conjecture 2.2.2 (Adjunction) *Let $(T/Z, B)$ be a lc pair of dim d such that $K_T + B \sim_{\mathbb{R}} 0/Z$. Define the unique class M_Z up to $\mathbb{R}$ -linear equivalence as $K_T + B \sim_{\mathbb{R}}^*(K_Z + D_Z + M_Z)$. Then the followings hold:*

Adjunction *We can choose an $M_Z \geq 0$ in its $\mathbb{R}$ -linear equivalence class such that $(Z, D_Z + M_Z)$ is lc.*

Effective adjunction *Fix Γ_f . Then there is a constant $I \in \mathbb{N}$ depending only on d and Γ_f such that $|IM_Z|$ is a free linear system for an appropriate choice of M_Z . In addition the following holds*

$$I(K_T + B) \sim^* I(K_Z + D_Z + M_Z).$$

It is expected that the effective adjunction implies the boundedness of S_1 under our assumptions.

8. If the third case occurs in the division above then we need to repeat the pprocess with a bigger ϵ . We have new coefficients in the boundary. Moreover we need to prove that this process stops after a bounded number of steps.
9. If every time the third case happens then at the end we get a pair (X_r, B_r) which is terminal in codim 2 and $-(K_{X_r} + B_r)$ is nef and big. After some more blow ups and blow downs we may prove that the index of $K_{X_r} + B_r$ is bounded.

3 List of notation and terminology

$\mathbb{N}$	<i>The set of natural numbers $\{1, 2, \dots\}$.</i>
$\mathbb{R}^+$	<i>The set of positive real numbers. Similar notation for $\mathbb{Q}$.</i>
dim	<i>Dimension or dimensional.</i>
WLF	<i>Weak log Fano. $(X/Z, B)$ is WLF if X/Z is a projective contraction and $-(K_X + B)$ is nef and big/Z and X is $\mathbb{Q}$-factorial.</i>
Pseudo-WLF	<i>Pseudo weak log Fano/Z i.e. there is a B where $(X/Z, B)$ is WLF.</i>
Φ_{sm}	<i>The set of standard boundary coefficients i.e. $\{\frac{k-1}{k}\}_{k \in \mathbb{N} \cup \{1\}}$.</i>
$\Gamma_{\mathbf{f}}$	<i>A finite subset of $[0, 1]$.</i>
$\text{mld}(\mu, \mathbf{X}, \mathbf{B})$	<i>The log minimal discrepancy of (X, B) at the centre μ.</i>
$\text{index}_{\mathbf{P}}(\mathbf{D})$	<i>The smallest positive natural number r s.t. rD is a Cartier divisor at P.</i>
$\text{WC}_{\delta, \mathbf{d}, \Gamma}$	<i>The weak conjecture on the boundedness of ϵ-lc complements in dim d. See 1.1.2</i>
$\text{SC}_{\delta, \mathbf{d}}$	<i>The strong conjecture on the boundedness of ϵ-lc complements in dim d. See 1.1.3</i>
$\text{BAB}_{\delta, \mathbf{d}, \Gamma}$	<i>The Alexeev-Borisovs conjecture on the boundedness of d-dim δ-lc WLF varieties. See 1.1.4</i>
$\text{LT}_{\mathbf{d}}$	<i>The log termination conjecture in dim d. See 1.1.7</i>
$\text{ACC}_{\mathbf{d}, \Gamma}$	<i>The ACC conjecture on mlds in dim d. See 1.1.6</i>

4 References:

- [A1] V. Alexeev; *Boundedness and K^2 for log surfaces*. Internat. J. Math. 5 (1994), no. 6, 779–810.
- [A2] V. Alexeev; *Two two dimensional terminations*. Duke Math. J. 69 (1993), no. 3, 527–545.
- [AM] V. Alexeev, S. Mori; *Bounding singular surfaces of general type*. Algebra, arithmetic and geometry with applications (West Lafayette, IN, 2000), 143–174, Springer, Berlin, 2004.
- [Am] F. Ambro; *On minimal log discrepancies*. Math. Res. Lett. 6 (1999), no. 5-6, 573–580.
- [B] A.A. Borisov; *Boundedness of Fano threefolds with log-terminal singularities of given index*. J. Math. Sci. Univ. Tokyo 8 (2001), no. 2, 329–342.
- [C] A. Corti; *Recent results in higher-dimensional birational geometry*. Current topics in complex algebraic geometry (Berkeley, CA, 1992/93), 35–56, Math. Sci. Res. Inst. Publ., 28, Cambridge Univ. Press, Cambridge, 1995.
- [C1] A. Corti; *Singularities of linear systems and 3-fold birational geometry*. Explicit birational geometry of 3-folds, 259–312, London Math. Soc. Lecture Note Ser., 281, Cambridge Univ. Press, Cambridge, 2000.
- [CR] A. Corti, M. Reid; *Explicit birational geometry of 3-folds*. Edited by Alessio Corti and Miles Reid. London Mathematical Society Lecture Note Series, 281. Cambridge University Press, Cambridge, 2000.
- [CPR] A. Corti, A. Pukhlikov, M. Reid; *Fano 3-fold hypersurfaces*. Explicit birational geometry of 3-folds, 175–258, London Math. Soc. Lecture Note Ser., 281, Cambridge Univ. Press, Cambridge, 2000.

- [H] R. Hartshorne; *Algebraic geometry*. Graduate Texts in Mathematics, No. 52. Springer-Verlag, 1977.
- [K1] Y. Kawamata; *Boundedness of $\mathbf{Q}$ -Fano threefolds*. Proceedings of the International Conference on Algebra, Part 3 (Novosibirsk, 1989), 439–445, Contemp. Math., 131, Part 3, Amer. Math. Soc., Providence, RI, 1992.
- [K2] Y. Kawamata; *Termination of log flips for algebraic 3-folds*. Internat. J. Math. 3 (1992), no. 5, 653–659.
- [K3] Y. Kawamata; *Subadjunction of log canonical divisors for a subvariety of codimension 2*. Birational algebraic geometry (Baltimore, MD, 1996), 79–88, Contemp. Math., 207, Amer. Math. Soc., Providence, RI, 1997.
- [K4] Y. Kawamata; *Subadjunction of log canonical divisors. II*. Amer. J. Math. 120 (1998), no. 5, 893–899.
- [K5] Y. Kawamata; *The number of the minimal models for a 3-fold of general type is finite*. Math. Ann. 276 (1987), no. 4, 595–598.
- [K6] Y. Kawamata; *Termination of log flips in dimension 4*. Preprint. It contained a proof which turned to be not correct.
- [KMM] Y. Kawamata, K. Matsuda, K. Matsuki; *Introduction to the minimal model problem*. Algebraic geometry, Sendai, 1985, 283–360, Adv. Stud. Pure Math., 10, North-Holland, Amsterdam, 1987.
- [Ko1] J. Kollar; *Singularities of pairs*. Algebraic geometry—Santa Cruz 1995, 221–287, Proc. Sympos. Pure Math., 62, Part 1, Amer. Math. Soc., Providence, RI, 1997.

- [Ko2] J. Kollar; *Rational curves on algebraic varieties*. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], 32. Springer-Verlag, Berlin, 1996.
- [KD] J. Demailly, J. Kollar; *Semi-continuity of complex singularity exponents and Kähler-Einstein metrics on Fano orbifolds*. Ann. Sci. cole Norm. Sup. (4) 34 (2001), no. 4, 525–556.
- [KM] J. Kollar, S. Mori; *Birational geometry of algebraic varieties*. With the collaboration of C. H. Clemens and A. Corti. Translated from the 1998 Japanese original. Cambridge Tracts in Mathematics, 134. Cambridge University Press, Cambridge, 1998.
- [KMMT] J. Kollár, Y. Miyaoka, S. Mori, H. Takagi; *Boundedness of canonical $\mathbf{Q}$ -Fano 3-folds*. Proc. Japan Acad. Ser. A Math. Sci. 76 (2000), no. 5, 73–77.
- [K⁺] J. Kollár and others; *Flips and abundance for algebraic threefolds*. Papers from the Second Summer Seminar on Algebraic Geometry held at the University of Utah, Salt Lake City, Utah, August 1991. Astrisque No. 211 (1992). Socit Mathématique de France, Paris, 1992. pp. 1–258.
- [Mc] J. McKernan; *Boundedness of log terminal Fano pairs of bounded index*. ArXiv/math.AG/0205214
- [MP] J. McKernan, Yu. Prokhorov; *Threefold Thresholds*. ArXiv/math.AG/0205214
- [Pr] Yu. Prokhorov; *Lectures on complements on log surfaces*. MSJ Memoirs, 10. Mathematical Society of Japan, Tokyo, 2001.
- [Pr1] Yu. Prokhorov; *Boundedness of exceptional quotient singularities*. (Russian) Mat. Zametki 68 (2000), no. 5, 786–789; translation in Math. Notes 68 (2000), no. 5-6, 664–667

- [Pr2] Yu. Prokhorov; *Boundedness of nonbirational extremal contractions*. Internat. J. Math. 11 (2000), no. 3, 393–411.
- [PrM] D. Markushevich, Yu. Prokhorov; *Exceptional quotient singularities*. Amer. J. Math. 121 (1999), no. 6, 1179–1189.
- [PrI] V.A. Iskovskikh, Yu. Prokhorov; *Fano varieties*. Algebraic geometry, V, 1–247, *Encyclopaedia Math. Sci.*, 47, Springer, Berlin, 1999.
- [PSh] Yu. Prokhorov; V.V. Shokurov; *The first fundamental theorem on complements: from global to local*. (Russian) Izv. Ross. Akad. Nauk Ser. Mat. 65 (2001), no. 6, 99–128; translation in Izv. Math. 65 (2001), no. 6, 1169–1196.
- [PSh1] Yu. Prokhorov; V.V. Shokurov; *Toward the second main theorem on complements: from local to global*. Preprint 2001.
- [R] M. Reid; *Chapters on algebraic surfaces*. Complex algebraic geometry (Park City, UT, 1993), 3–159, IAS/Park City Math. Ser., 3, Amer. Math. Soc., Providence, RI, 1997.
- [R1] M. Reid; *Update on 3-folds*. Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002), 513–524, Higher Ed. Press, Beijing, 2002.
- [R2] M. Reid; *Twenty-five years of 3-folds—an old person’s view*. Explicit birational geometry of 3-folds, 313–343, London Math. Soc. Lecture Note Ser., 281, Cambridge Univ. Press, Cambridge, 2000.
- [R3] M. Reid; *Young person’s guide to canonical singularities*. Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), 345–414, Proc. Sympos. Pure Math., 46, Part 1, Amer. Math. Soc., Providence, RI, 1987.
- [R4] M. Reid; *The moduli space of 3-folds with $K = 0$ may nevertheless be irreducible*. Math. Ann. 278 (1987), no. 1–4, 329–334.

- [Sh1] V.V. Shokurov; *Three-dimensional log flips*. With an appendix in English by Yujiro Kawamata. Russian Acad. Sci. Izv. Math. 40 (1993), no. 1, 95–202.
- [Sh2] V.V. Shokurov; *Complements on surfaces*. Algebraic geometry, 10. J. Math. Sci. (New York) 102 (2000), no. 2, 3876–3932.
- [Sh3] V.V. Shokurov; *Prelimiting flips*. Tr. Mat. Inst. Steklova 240 (2003), Biratsion. Geom. Linein. Sist. Konechno Porozhdennye Algebry, 82–219; translation in Proc. Steklov Inst. Math. 2003, no. 1 (240), 75–213.
- [Sh4] V.V. Shokurov; *Letters of a birationalist V: Mld's and termination of log flips*.
- [Sh5] V.V. Shokurov; *3-fold log models*. Algebraic geometry, 4. J. Math. Sci. 81 (1996), no. 3, 2667–2699.
- [Sh6] V.V. Shokurov; *Letters of a bi-rationalist. IV. Geometry of log flips*. Algebraic geometry, 313–328, de Gruyter, Berlin, 2002.
- [Sh7] V.V. Shokurov; *A nonvanishing theorem*. (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 49 (1985), no. 3, 635–651.
- [Sh8] V.V. Shokurov; *ACC in codim 2*. Preprint.